

Mathematical Methods - Mathematical Analysis: takehome (simulation)

Suppose $T : \ell^2(\mathbb{Z}) \rightarrow \ell^2(\mathbb{Z})$ is defined as

$$\widehat{Ta}(t) = \begin{cases} 0 & \text{if } -\pi \leq t \leq 0, \\ \left(\frac{\pi}{2} - \left|t - \frac{\pi}{2}\right|\right) \cdot \hat{a}(t) & \text{if } 0 < t \leq \pi. \end{cases}$$

for all $a \in \ell^2(\mathbb{Z})$.

- Is T linear?
- Is T stable?
- Is T time-invariant?
- Is T realizable?

Motivate your answer.

Find $b \in \ell^2(\mathbb{Z})$ such that

$$Ta(n) = \sum_{k=-\infty}^{+\infty} a(n-k)b(k).$$

Compute $T\delta_b$, where b is the last digit of your student ID.

It is useful to know that:

- $\hat{a}(t) = \sum_{k=-\infty}^{+\infty} a(k)e^{-ikt}$;
- $\delta_k : \mathbb{Z} \rightarrow \mathbb{C}$ is a discrete Dirac's delta, defined as

$$\delta_k(n) = \begin{cases} 1 & \text{if } n = k, \\ \hat{a}(t) & \text{if } n \neq k. \end{cases}$$

- for the meaning of linear, stable, time-invariant and realizable see your notes from the class, or the solution to the exercise.