

On the Eigenvalues of a Class of Saddle Point Matrices

V. Simoncini



*Dipartimento di Matematica
Università di Bologna
valeria@dm.unibo.it*

Thanks to

Mario Arioli, *RAL, UK*

Michele Benzi, *Emory University (GA)*

Ilaria Perugia, *Università di Pavia*

Miro Rozložník, *Institute of Computer Science, Academy of
Science, Prague*

Application problems

- Computational Fluid Dynamics
- Elasticity problems
- Mixed (FE) formulations of II and IV order elliptic PDEs
- Linearly Constrained Programs
- Linear Regression in Statistics
- Weighted Least Squares (Image restoration)
- ... Survey: Benzi, Golub and Liesen, Acta Num 2005

$$\begin{bmatrix} A & B^T \\ B & -C \end{bmatrix} \begin{bmatrix} u \\ v \end{bmatrix} = \begin{bmatrix} f \\ g \end{bmatrix}$$

Motivational Application

Constrained Quadratic minimization problem

$$\text{minimize } J(u) = \frac{1}{2} \langle Au, u \rangle - \langle f, u \rangle$$

subject to $Bu = g$

A $n \times n$ spd, B $m \times n$, $m \leq n$ full rank



Lagrange multipliers approach

Karush-Kuhn-Tucker (KKT) system

Spectral properties

- $\mathcal{M} = \begin{bmatrix} A & B^T \\ B & O \end{bmatrix}$ $0 < \lambda_n \leq \dots \leq \lambda_1$ eigs of A
 $0 < \sigma_m \leq \dots \leq \sigma_1$ sing. vals of B

Spectral properties

- $\mathcal{M} = \begin{bmatrix} A & B^T \\ B & O \end{bmatrix}$ $0 < \lambda_n \leq \dots \leq \lambda_1$ eigs of A
 $0 < \sigma_m \leq \dots \leq \sigma_1$ sing. vals of B

- (Rusten & Winther 1992) $\Lambda(\mathcal{M})$ subset of

$$\left[\frac{1}{2}(\lambda_n - \sqrt{\lambda_n^2 + 4\sigma_1^2}), \frac{1}{2}(\lambda_1 - \sqrt{\lambda_1^2 + 4\sigma_m^2}) \right] \cup \left[\lambda_n, \frac{1}{2}(\lambda_1 + \sqrt{\lambda_1^2 + 4\sigma_m^2}) \right]$$

Spectral properties

- $\mathcal{M} = \begin{bmatrix} A & B^T \\ B & -C \end{bmatrix}$ $0 < \lambda_n \leq \dots \leq \lambda_1$ eigs of A
 $0 < \sigma_m \leq \dots \leq \sigma_1$ sing. vals of B

- (Rusten & Winther 1992) $\Lambda(\mathcal{M})$ subset of

$$\left[\frac{1}{2}(\lambda_n - \sqrt{\lambda_n^2 + 4\sigma_1^2}), \frac{1}{2}(\lambda_1 - \sqrt{\lambda_1^2 + 4\sigma_m^2}) \right] \cup \left[\lambda_n, \frac{1}{2}(\lambda_1 + \sqrt{\lambda_1^2 + 4\sigma_m^2}) \right]$$

- (Silvester & Wathen 1994), $0 \leq \sigma_m \leq \dots \leq \sigma_1$

$$\lambda_n - \lambda_{\max}(C) - \sqrt{(\lambda_n + \lambda_{\max}(C))^2 + 4\sigma_1^2}$$

Spectral properties

- $\mathcal{M} = \begin{bmatrix} A & B^T \\ B & -C \end{bmatrix}$ $0 < \lambda_n \leq \dots \leq \lambda_1$ eigs of A
 $0 < \sigma_m \leq \dots \leq \sigma_1$ sing. vals of B
- (Rusten & Winther 1992) $\Lambda(\mathcal{M})$ subset of
$$\left[\frac{1}{2}(\lambda_n - \sqrt{\lambda_n^2 + 4\sigma_1^2}), \frac{1}{2}(\lambda_1 - \sqrt{\lambda_1^2 + 4\sigma_m^2}) \right] \cup \left[\lambda_n, \frac{1}{2}(\lambda_1 + \sqrt{\lambda_1^2 + 4\sigma_m^2}) \right]$$
- (Silvester & Wathen 1994), $0 \leq \sigma_m \leq \dots \leq \sigma_1$

$$\lambda_n - \lambda_{\max}(C) - \sqrt{(\lambda_n + \lambda_{\max}(C))^2 + 4\sigma_1^2}$$

More results for special cases (e.g., Perugia & S. 2000)

Block diagonal Preconditioner

$$\mathcal{P} = \begin{bmatrix} \tilde{A} & 0 \\ 0 & \tilde{C} \end{bmatrix} \quad \text{sym. pos. def.}$$

$$\tilde{A} \approx A \quad \tilde{C} \approx BA^{-1}B^T + C$$

Rusten Winther (1992), Silvester Wathen (1993-1994), Klawonn (1998)

Fischer Ramage Silvester Wathen (1998...), . . .

Block diagonal Preconditioner

$$\mathcal{P} = \begin{bmatrix} \tilde{A} & 0 \\ 0 & \tilde{C} \end{bmatrix} \quad \text{sym. pos. def.}$$

$$\tilde{A} \approx A \quad \tilde{C} \approx BA^{-1}B^T + C$$

Rusten Winther (1992), Silvester Wathen (1993-1994), Klawonn (1998)

Fischer Ramage Silvester Wathen (1998...), . . .

$\lambda \neq 0$ eigs of $\mathcal{P}^{-\frac{1}{2}} \mathcal{M} \mathcal{P}^{-\frac{1}{2}}$,

$$\lambda \in [-a, -b] \cup [c, d], \quad a, b, c, d > 0$$

Triangular preconditioner

$$A \text{ spd}, \quad \mathcal{P} = \begin{bmatrix} \tilde{A} & B^T \\ 0 & -\tilde{C} \end{bmatrix}$$

Bramble & Pasciak, Elman, Klawonn, Axelsson & Neytcheva, S. 2004

Triangular preconditioner

$$A \text{ spd}, \quad \mathcal{P} = \begin{bmatrix} \tilde{A} & B^T \\ 0 & -\tilde{C} \end{bmatrix}$$

Bramble & Pasciak, Elman, Klawonn, Axelsson & Neytcheva, S. 2004

Spectrum of $\mathcal{M}\mathcal{P}^{-1}$:

small complex cluster around 1 + real interval

Triangular preconditioner

$$A \text{ spd}, \quad \mathcal{P} = \begin{bmatrix} \tilde{A} & B^T \\ 0 & -\tilde{C} \end{bmatrix}$$

Bramble & Pasciak, Elman, Klawonn, Axelsson & Neytcheva, S. 2004

Spectrum of $\mathcal{M}\mathcal{P}^{-1}$:

small complex cluster around 1 + real interval

More precisely: $\theta \in \Lambda(\mathcal{M}\mathcal{P}^{-1})$

$$\Im(\theta) \neq 0 \Rightarrow |\theta - 1| \leq \sqrt{1 - \lambda_{\min}(A\tilde{A}^{-1})} \quad (\text{if } 1 - \lambda_{\min}(A\tilde{A}^{-1}) \geq 0)$$

$$\Im(\theta) = 0 \Rightarrow \theta \in [\chi_1, \chi_2] \ni 1$$

Constraint Preconditioner

$$Q = \begin{bmatrix} \tilde{A} & B^T \\ B & -C \end{bmatrix}$$

Axelsson (1979), Ewing Lazarov Lu Vassilevski (1990), ... many more papers 1997 -

Constraint Preconditioner

$$Q = \begin{bmatrix} \tilde{A} & B^T \\ B & -C \end{bmatrix}$$

Axelsson (1979), Ewing Lazarov Lu Vassilevski (1990), ... many more papers 1997 -

$\lambda \neq 0$ eigs of $\mathcal{M}Q^{-1}$, $\lambda \in \mathbb{R}^+$, $\lambda \in \{1\} \cup [\alpha_0, \alpha_1]$

Constraint Preconditioner

$$Q = \begin{bmatrix} \tilde{A} & B^T \\ B & -C \end{bmatrix}$$

Axelsson (1979), Ewing Lazarov Lu Vassilevski (1990), ... many more papers 1997 -

$\lambda \neq 0$ eigs of $\mathcal{M}Q^{-1}$, $\lambda \in \mathbb{R}^+$, $\lambda \in \{1\} \cup [\alpha_0, \alpha_1]$

Remark: $Bx_k = 0$ for all iterates x_k ($C = 0$)

Constraint Preconditioner

$$Q = \begin{bmatrix} \tilde{A} & B^T \\ B & -C \end{bmatrix}$$

Axelsson (1979), Ewing Lazarov Lu Vassilevski (1990), ... many more papers 1997 -

$\lambda \neq 0$ eigs of $\mathcal{M}Q^{-1}$, $\lambda \in \mathbb{R}^+$, $\lambda \in \{1\} \cup [\alpha_0, \alpha_1]$

Remark: $Bx_k = 0$ for all iterates x_k ($C = 0$)

$$Q^{-1} = \begin{bmatrix} I & -B^T \\ O & I \end{bmatrix} \begin{bmatrix} I & O \\ O & -(\mathbf{B}\mathbf{B}^T + \mathbf{C})^{-1} \end{bmatrix} \begin{bmatrix} I & O \\ -B & I \end{bmatrix}$$

($\tilde{A} = I$ if prescaling used)

Computational Considerations.

3D Magnetostatic problem

$H \approx BB^T + C$ with H : Incomplete Cholesky fact.

(ICT package, Saad & Chow)

Computational Considerations.

3D Magnetostatic problem

$$H \approx BB^T + C \quad \text{with } H: \text{ Incomplete Cholesky fact.}$$

(ICT package, Saad & Chow)

Elapsed Time

size	MA27	QMR		ILDLT(10)
		Q	$\hat{Q}(2)(it)$	
1119	0.6	3.0	1.7(18)	0.7
2208	2.3	11.7	3.1(18)	1.5
4371	10.2	64.6	8.4(20)	5.2
8622	83.4	466.0	18.3(29)	31.0
22675	753.5	3745.5	63.2(45)	246.0

Computational Considerations.

3D Magnetostatic problem

$$H \approx BB^T + C \quad \text{with } H: \text{ Incomplete Cholesky fact.}$$

(ICT package, Saad & Chow)


Elapsed Time

size	MA27	QMR		QMR ILDLT(10)	QMR	
		Q	$\hat{Q}(2)(it)$		\mathcal{P}	$\hat{\mathcal{P}}(2)(it)$
1119	0.6	3.0	1.7(18)	0.7	4.0	2.0
2208	2.3	11.7	3.1(18)	1.5	15.2	4.9
4371	10.2	64.6	8.4(20)	5.2	73.9	11.0
8622	83.4	466.0	18.3(29)	31.0	510.1	24.3
22675	753.5	3745.5	63.2(45)	246.0	4161.4	128.2

Spectrum of perturbed problem





3D Magnetostatic problem



Spectrum of perturbed problem

3D Magnetostatic problem



$$\|(BB^T + C) - H\|_{\infty} \approx 2 \cdot 10^{-1} \|BB^T + C\|_{\infty}$$

Eigenvalue problem

$$\hat{Q} = \begin{bmatrix} I & O \\ B & I \end{bmatrix} \begin{bmatrix} I & O \\ O & -H \end{bmatrix} \begin{bmatrix} I & B^T \\ O & I \end{bmatrix}, \quad H \approx BB^T, \quad C = O$$

$$\begin{bmatrix} A & B^T \\ B & O \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \lambda \hat{Q} \begin{bmatrix} x \\ y \end{bmatrix}$$

Eigenvalue problem

$$\hat{Q} = \begin{bmatrix} I & O \\ B & I \end{bmatrix} \begin{bmatrix} I & O \\ O & -H \end{bmatrix} \begin{bmatrix} I & B^T \\ O & I \end{bmatrix}, \quad H \approx BB^T, \quad C = O$$

$$\begin{bmatrix} A & B^T \\ B & O \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \lambda \hat{Q} \begin{bmatrix} x \\ y \end{bmatrix}$$

can be written as

$$\begin{bmatrix} I & O \\ -B & I \end{bmatrix} \begin{bmatrix} A & B^T \\ B & O \end{bmatrix} \begin{bmatrix} I & -B^T \\ O & I \end{bmatrix} \begin{bmatrix} u \\ v \end{bmatrix} = \lambda \begin{bmatrix} I & O \\ O & -H \end{bmatrix} \begin{bmatrix} u \\ v \end{bmatrix}$$

$$\begin{bmatrix} A & (I - A)B^T \\ B(I - A) & -B(2I - A)B^T \end{bmatrix} \begin{bmatrix} u \\ v \end{bmatrix} = \lambda \begin{bmatrix} I & O \\ O & -H \end{bmatrix} \begin{bmatrix} u \\ v \end{bmatrix}$$

A “different” perspective

$$\begin{bmatrix} A & B^T \\ -B & C \end{bmatrix} \begin{bmatrix} u \\ v \end{bmatrix} = \begin{bmatrix} f \\ -g \end{bmatrix} \quad \mathcal{M}_- x = d$$

Polyak 1970, ..., Wathen & Fischer & Silvester 1995, Fischer & Ramage & Silvester & Wathen 1998, Fischer & Peherstorfer 2001, Bai & Golub & Ng 2003, Sidi 2003, Benzi & Gander & Golub 2003, Benzi & Golub 2004, S. & Benzi 2004, ...

A “different” perspective

$$\begin{bmatrix} A & B^T \\ -B & C \end{bmatrix} \begin{bmatrix} u \\ v \end{bmatrix} = \begin{bmatrix} f \\ -g \end{bmatrix} \quad \mathcal{M}_- x = d$$

Polyak 1970, ..., Wathen & Fischer & Silvester 1995, Fischer & Ramage & Silvester & Wathen 1998, Fischer & Peherstorfer 2001, Bai & Golub & Ng 2003, Sidi 2003, Benzi & Gander & Golub 2003, Benzi & Golub 2004, S. & Benzi 2004, ...

- \mathcal{M} positive real $\Rightarrow \Lambda(\mathcal{M}_-) \text{ in } \mathbb{C}^+$

A “different” perspective

$$\begin{bmatrix} A & B^T \\ -B & C \end{bmatrix} \begin{bmatrix} u \\ v \end{bmatrix} = \begin{bmatrix} f \\ -g \end{bmatrix} \quad \mathcal{M}_- x = d$$

Polyak 1970, ..., Wathen & Fischer & Silvester 1995, Fischer & Ramage & Silvester & Wathen 1998, Fischer & Peherstorfer 2001, Bai & Golub & Ng 2003, Sidi 2003, Benzi & Gander & Golub 2003, Benzi & Golub 2004, S. & Benzi 2004, ...

- \mathcal{M} positive real $\Rightarrow \Lambda(\mathcal{M}_-) \text{ in } \mathbb{C}^+$
- More refined spectral information possible

A “different” perspective

$$\begin{bmatrix} A & B^T \\ -B & C \end{bmatrix} \begin{bmatrix} u \\ v \end{bmatrix} = \begin{bmatrix} f \\ -g \end{bmatrix} \quad \mathcal{M}_- x = d$$

Polyak 1970, ..., Wathen & Fischer & Silvester 1995, Fischer & Ramage & Silvester & Wathen 1998, Fischer & Peherstorfer 2001, Bai & Golub & Ng 2003, Sidi 2003, Benzi & Gander & Golub 2003, Benzi & Golub 2004, S. & Benzi 2004, ...

- \mathcal{M} positive real $\Rightarrow \Lambda(\mathcal{M}_-) \text{ in } \mathbb{C}^+$
- More refined spectral information possible
- New classes of preconditioners

A “different” perspective

$$\begin{bmatrix} A & B^T \\ -B & C \end{bmatrix} \begin{bmatrix} u \\ v \end{bmatrix} = \begin{bmatrix} f \\ -g \end{bmatrix} \quad \mathcal{M}_- x = d$$

Polyak 1970, ..., Wathen & Fischer & Silvester 1995, Fischer & Ramage & Silvester & Wathen 1998, Fischer & Peherstorfer 2001, Bai & Golub & Ng 2003, Sidi 2003, Benzi & Gander & Golub 2003, Benzi & Golub 2004, S. & Benzi 2004, ...

- \mathcal{M} positive real $\Rightarrow \Lambda(\mathcal{M}_-) \text{ in } \mathbb{C}^+$
- More refined spectral information possible
- New classes of preconditioners
- General framework for spectral analysis of some indefinite preconditioners

Benzi & Simoncini (tr 2005)

Spectral properties of \mathcal{M}_-

$$\mathcal{M}_- = \begin{bmatrix} A & B^T \\ -B & C \end{bmatrix}$$

A $n \times n$ sym. semidef. matrix, B $m \times n$, $m \leq n$

\mathcal{M}_- has at least $n - m$ real eigenvalues

Detailed analysis for $A = \eta I, C = O$ in:

- ★ Fischer & Ramage & Silvester & Wathen 1998
- ★ Fischer & Peherstorfer 2001

Reality condition

Let A be spd and $C = \beta I$. If

$$\lambda_{\min}(A + \beta I) \geq 4 \lambda_{\max}(B^T A^{-1} B + \beta I),$$

then all eigenvalues of \mathcal{M}_- are real (and positive)

Reality condition

Let A be spd and $C = \beta I$. If

$$\lambda_{\min}(A + \beta I) \geq 4 \lambda_{\max}(B^T A^{-1} B + \beta I),$$

then all eigenvalues of \mathcal{M}_- are real (and positive)

The condition is sufficient, but not necessary:

$$\mathcal{M}_- = \left[\begin{array}{cc|c} \frac{1}{2} & 0 & 0 \\ 0 & 3 & 1 \\ \hline 0 & -1 & 0 \end{array} \right], \quad \begin{aligned} \lambda_{\min}(A) &= \frac{1}{2} \\ \lambda_{\max}(B A^{-1} B) &= \frac{1}{3} \end{aligned}$$

has real spectrum, but does not satisfy the condition

The (steady-state) Stokes problem

$$\left\{ \begin{array}{ll} -\Delta \mathbf{u} + \nabla p = \mathbf{f} & \text{in } \Omega \\ \nabla \cdot \mathbf{u} = 0 & \text{in } \Omega \\ \mathcal{B}\mathbf{u} = \mathbf{g} & \text{on } \Gamma. \end{array} \right. \Rightarrow \mathcal{M}_{\pm}$$

The (steady-state) Stokes problem

$$\left\{ \begin{array}{ll} -\Delta \mathbf{u} + \nabla p = \mathbf{f} & \text{in } \Omega \\ \nabla \cdot \mathbf{u} = 0 & \text{in } \Omega \\ \mathcal{B}\mathbf{u} = \mathbf{g} & \text{on } \Gamma. \end{array} \right. \Rightarrow \mathcal{M}_{\pm}$$

Numerical evidence: eigs of \mathcal{M}_- are **all** real and positive

(for several discretizations and b.c.)

The (steady-state) Stokes problem

$$\begin{cases} -\Delta \mathbf{u} + \nabla p = \mathbf{f} & \text{in } \Omega \\ \nabla \cdot \mathbf{u} = 0 & \text{in } \Omega \\ \mathcal{B}\mathbf{u} = \mathbf{g} & \text{on } \Gamma. \end{cases} \Rightarrow \mathcal{M}_{\pm}$$

Numerical evidence: eigs of \mathcal{M}_- are **all** real and positive

(for several discretizations and b.c.)

Explanation:

$\Omega = [0, 1] \times [0, 1]$, Dirichlet b.c. div-stable discr.

$$\begin{aligned} \lambda_{\min}(A) &\approx 2\pi^2 \approx 19, \\ \lambda_{\max}(BA^{-1}B^T) &\approx 1 \end{aligned} \Rightarrow \lambda_{\min}(A) > 4\lambda_{\max}(BA^{-1}B^T)$$

(In)definite inner product

$$\mathcal{M}_- = \begin{bmatrix} A & B^T \\ -B & C \end{bmatrix}$$

$$J = \begin{bmatrix} I & O \\ O & -I \end{bmatrix}$$

J indefinite

\mathcal{M}_- is J -sym.

(In)definite inner product

$$\mathcal{M}_- = \begin{bmatrix} A & B^T \\ -B & C \end{bmatrix} \quad J = \begin{bmatrix} I & O \\ O & -I \end{bmatrix} \quad \begin{array}{l} J \text{ indefinite} \\ \mathcal{M}_- \text{ is } J\text{-sym.} \end{array}$$

If $\Lambda(\mathcal{M}_-) \subset \mathbb{R}^+$, there exists a nonstandard inner product with which \mathcal{M}_- is spd (and diag.ble)

(In)definite inner product

$$\mathcal{M}_- = \begin{bmatrix} A & B^T \\ -B & C \end{bmatrix} \quad J = \begin{bmatrix} I & O \\ O & -I \end{bmatrix} \quad \begin{array}{l} J \text{ indefinite} \\ \mathcal{M}_- \text{ is } J\text{-sym.} \end{array}$$

If $\Lambda(\mathcal{M}_-) \subset \mathbb{R}^+$, there exists a nonstandard inner product with which \mathcal{M}_- is spd (and diag.ble)

$$G = \begin{bmatrix} A - \gamma I & B^T \\ B & \gamma I \end{bmatrix}, \quad \mathcal{M}_- G = G \mathcal{M}_-^T$$

(In)definite inner product

$$\mathcal{M}_- = \begin{bmatrix} A & B^T \\ -B & C \end{bmatrix} \quad J = \begin{bmatrix} I & O \\ O & -I \end{bmatrix} \quad \begin{array}{l} J \text{ indefinite} \\ \mathcal{M}_- \text{ is } J\text{-sym.} \end{array}$$

If $\Lambda(\mathcal{M}_-) \subset \mathbb{R}^+$, there exists a nonstandard inner product with which \mathcal{M}_- is spd (and diag.ble)

$$G = \begin{bmatrix} A - \gamma I & B^T \\ B & \gamma I \end{bmatrix}, \quad \mathcal{M}_- G = G \mathcal{M}_-^T$$

If $\gamma = \frac{1}{2} \lambda_{\min}(A)$, $\lambda_{\min}(A) > 4\lambda_{\max}(BA^{-1}B^T)$ (with $C = O$)

$\Rightarrow G$ is spd $\Rightarrow G$ defines an inner product for \mathcal{M}_-

$$\mathcal{M}_- = \begin{bmatrix} A & B^T \\ -B & C \end{bmatrix} = \begin{bmatrix} A & O \\ O & C \end{bmatrix} + \begin{bmatrix} O & B^T \\ -B & O \end{bmatrix}$$
$$= \mathcal{H} + \mathcal{S}$$

$$\mathcal{M}_- = \begin{bmatrix} A & B^T \\ -B & C \end{bmatrix} = \begin{bmatrix} A & O \\ O & C \end{bmatrix} + \begin{bmatrix} O & B^T \\ -B & O \end{bmatrix}$$
$$= \mathcal{H} + \mathcal{S}$$

Use the preconditioner

$$\mathcal{R}_\alpha = \frac{1}{2\alpha} (\mathcal{H} + \alpha I)(\mathcal{S} + \alpha I) \quad \alpha \in \mathbb{R}, \alpha > 0$$

(Bai & Golub & Ng 2003), Benzi & Gander & Golub 2003, Benzi & Golub 2004, S. & Benzi 2004, Benzi & Ng, 2004

Spectral bounds: *Simoncini & Benzi 2004*

Reality condition

Assume A is spd and $C = 0$. If

$$\alpha \leq \frac{1}{2} \lambda_{\min}(A)$$

then all eigenvalues of $\mathcal{R}_\alpha^{-1} \mathcal{M}_-$ are real (Simoncini & Benzi '04)

Reality condition

Assume A is spd and $C = 0$. If

$$\alpha \leq \frac{1}{2} \lambda_{\min}(A)$$

then all eigenvalues of $\mathcal{R}_\alpha^{-1} \mathcal{M}_-$ are real (Simoncini & Benzi '04)

But even more general:

If $\lambda_{\min}(A) > 4\lambda_{\max}(BA^{-1}B^T)$ then

all eigenvalues are **real** for **any** α

(cf. Stokes problem)

Location of \mathcal{M}_- 's spectrum

Let $\lambda \in \Lambda(\mathcal{M}_-)$, $x = [u; v]$ e'vect., A spd (cf. Sidi '03, $C = O$)

★ If $\Im(\lambda) \neq 0$ then

$$\begin{aligned} \frac{1}{2}(\lambda_{\min}(A) + \lambda_{\min}(C)) &\leq \Re(\lambda) \leq \frac{1}{2}(\lambda_{\max}(A) + \lambda_{\max}(C)) \\ |\Im(\lambda)| &\leq \sigma_{\max}(B). \end{aligned}$$

Location of \mathcal{M}_- 's spectrum

Let $\lambda \in \Lambda(\mathcal{M}_-)$, $x = [u; v]$ e'vect., A spd (cf. Sidi '03, $C = O$)

★ If $\Im(\lambda) \neq 0$ then

$$\begin{aligned} \frac{1}{2}(\lambda_{\min}(A) + \lambda_{\min}(C)) &\leq \Re(\lambda) \leq \frac{1}{2}(\lambda_{\max}(A) + \lambda_{\max}(C)) \\ |\Im(\lambda)| &\leq \sigma_{\max}(B). \end{aligned}$$

★ If $\Im(\lambda) = 0$ then for $v \neq 0$

$$2 \min\{\lambda_{\min}(A), \lambda_{\min}(C)\} \leq \lambda \leq \max\{\lambda_{\max}(A), \lambda_{\max}(C)\}$$

and for $v = 0$, $\lambda_{\min}(A) \leq \lambda \leq \lambda_{\max}(A)$

Location of \mathcal{M}_- 's spectrum

Let $\lambda \in \Lambda(\mathcal{M}_-)$, $x = [u; v]$ e'vect., A spd (cf. Sidi '03, $C = O$)

★ If $\Im(\lambda) \neq 0$ then

$$\frac{1}{2}(\lambda_{\min}(A) + \lambda_{\min}(C)) \leq \Re(\lambda) \leq \frac{1}{2}(\lambda_{\max}(A) + \lambda_{\max}(C))$$
$$|\Im(\lambda)| \leq \sigma_{\max}(B).$$

★ If $\Im(\lambda) = 0$ then for $v \neq 0$

$$2 \min\{\lambda_{\min}(A), \lambda_{\min}(C)\} \leq \lambda \leq \max\{\lambda_{\max}(A), \lambda_{\max}(C)\}$$

and for $v = 0$, $\lambda_{\min}(A) \leq \lambda \leq \lambda_{\max}(A)$

$$\mathcal{M}_- = \left[\begin{array}{cc|c} 2 & 0 & 1 \\ 0 & 1 & 0 \\ \hline -1 & 0 & 1 \end{array} \right] \quad \lambda_1 = 1, \quad \lambda_{2,3} = \frac{3}{2} \pm i \frac{\sqrt{3}}{2}.$$

Inexact Indefinite Precond.

$$\begin{bmatrix} A & (I - A)B^T \\ B(I - A) & -B(2I - A)B^T \end{bmatrix} \begin{bmatrix} u \\ v \end{bmatrix} = \lambda \begin{bmatrix} I & O \\ O & -H \end{bmatrix} \begin{bmatrix} u \\ v \end{bmatrix}$$

Inexact Indefinite Precond.

$$\begin{bmatrix} A & (I - A)B^T \\ B(I - A) & -B(2I - A)B^T \end{bmatrix} \begin{bmatrix} u \\ v \end{bmatrix} = \lambda \begin{bmatrix} I & O \\ O & -H \end{bmatrix} \begin{bmatrix} u \\ v \end{bmatrix}$$

Eigenvalue bounds: Let $\hat{C} = B(2I - A)B^T H^{-1}$


★ If $\Im(\lambda) \neq 0$ then

$$\begin{aligned} \frac{1}{2}(\lambda_{\min}(A) + \lambda_{\min}(\hat{C})) &\leq \Re(\lambda) \leq \frac{1}{2}(\lambda_{\max}(A) + \lambda_{\max}(\hat{C})) \\ |\Im(\lambda)| &\leq \sigma_{\max}((I - A)BH^{-\frac{1}{2}}). \end{aligned}$$

★ If $\Im(\lambda) = 0$ then

$$2 \min\{\lambda_{\min}(A), \lambda_{\min}(\hat{C})\} \leq \lambda \leq \max\{\lambda_{\max}(A), \lambda_{\max}(\hat{C})\}$$

Spectral bounds



Final Considerations

- New framework increases understanding

Final Considerations

- New framework increases understanding
- But: in general eigenvector analysis may be challenging

Final Considerations

- New framework increases understanding
- But: in general eigenvector analysis may be challenging
- General non-Hermitian problem not fully understood

Final Considerations

- New framework increases understanding
- But: in general eigenvector analysis may be challenging
- General non-Hermitian problem not fully understood
- Visit `http://www.dm.unibo.it/~simoncin`