



**Extended Krylov subspace method
for matrix functions: new theoretical and
computational results**

V. Simoncini

Dipartimento di Matematica, Università di Bologna

valeria@dm.unibo.it

Joint work in progress with Leonid Knizhnerman, Moscow

The problem

Given $A \in \mathbb{R}^{n \times n}$, $v \in \mathbb{R}^n$ and f sufficiently smooth function, approximate

$$x = f(A)v$$

★ A large dimensions, $\|v\| = 1$

Projection-type methods

\mathcal{K} approximation space, $m = \dim(\mathcal{K})$ $V \in \mathbb{R}^{n \times m}$ s.t. $\mathcal{K} = \text{range}(V)$

$$x = f(A)v \quad \approx \quad x_m = Vf(V^\top AV)(V^\top v)$$

Standard Krylov subspace approximation

$$\mathcal{K} = K_m(A, v)$$

For $H_m = V^\top AV$, $v = Ve_1$ and $V^\top V = I_m$:

$$x_m = V_m f(H_m) e_1$$

Polynomial approximation: $x_m = p_{m-1}(A)v$

(p_{m-1} interpolates f at eigenvalues of H_m)

Note: Procedure valid for A symmetric and nonsymmetric

★ Numerical and theoretical results since mid '80s.

Acceleration Procedures: Shift-Invert Lanczos

A symmetric pos. (semi)def.

Choose γ s.t. $(I + \gamma A)$ is invertible, and construct

$$\mathcal{K} = K_m((I + \gamma A)^{-1}, v), \quad \text{van den Eshof-Hochbruck '06, Moret-Novati '04}$$

with $T_m = V^\top (I + \gamma A)^{-1} V$, $v = V e_1$ and $V^\top V = I_m$

$$x_m = V_m f\left(\frac{1}{\gamma}(T_m^{-1} - I_m)\right)e_1$$

Rational approximation: $x_m = p_{m-1}((I + \gamma A)^{-1})v$

Choice of γ : $\gamma = 1/\sqrt{\lambda_{\min}\lambda_{\max}}$ (Moret, tr 2005)

Acceleration Procedures: Extended Krylov

For A nonsingular,

$$\mathcal{K} = K_{m_1}(A, v) + K_{m_2}(A^{-1}, v), \quad \text{Druskin-Knizhnerman 1998, } A \text{ sym.}$$

Note: $\mathcal{K} = A^{-(m_2-1)} K_{m_1+m_2-1}(A, v)$

Algorithm (augmentation-style)

- Fix $m_2 \ll m_1$
- Run m_2 steps of Inverted Lanczos
- Run m_1 steps of Standard Lanczos + orth.

Extended Krylov: a new implementation

$m_1 = m_2 = m$ not fixed a priori

$$\mathcal{K} = K_m(A, v) + K_m(A^{-1}, A^{-1}v)$$

★ Arnoldi-type recurrence:

- $U_1 \leftarrow [v, A^{-1}v] + \text{orth}$

- $U_{j+1} \leftarrow [AU_j(:, 1), A^{-1}U_j(:, 2)] + \text{orth} \quad j = 1, 2, \dots$

★ Recurrence to cheaply compute $\mathcal{T}_m = \mathcal{U}_m^\top A \mathcal{U}_m$, $\mathcal{U}_m = [U_1, \dots, U_m]$

★ Compute $x_m = \mathcal{U}_m f(\mathcal{T}_m) e_1$

Simoncini, 2007

Extended Krylov: Convergence theory I

f satisfying $f(z) = \int_{-\infty}^0 \frac{1}{z - \zeta} d\mu(\zeta), \quad z \in \mathbb{C} \setminus]-\infty, 0]$

(with convenient measure $d\mu(\zeta)$)

Druskin-Knizhnerman 1998:

A sym: $\|x - x_m\| = \mathcal{O}(m^2 \exp(-2m \sqrt[4]{\frac{\lambda_{\min}}{\lambda_{\max}}}))$

Extended Krylov: Convergence theory II

A new approximation result for nonsingular A :

Let $f = f_1 + f_2$, $a \in [0, \infty)$

$$\|f_1(z) - \sum_{k=0}^{m-1} \gamma_{1,k} F_{1,k}(z)\| \leq c_1 \Phi_1(-a)^{-m},$$

$$\|f_2(z) - \sum_{k=0}^{m-1} \gamma_{2,k} F_{2,k}(z^{-1})\| \leq c_2 \Phi_2(-\frac{1}{a})^{-m},$$


$\Phi_1, F_{1,k}$ conformal mapping and Faber Polynomial w.r.to $W(A)$

$\Phi_2, F_{2,k}$ conformal mapping and Faber Polynomial w.r.to $W(A^{-1})$

From this, for A symmetric: $\|x - x_m\| = O(\exp(-2m \sqrt[4]{\frac{\lambda_{\min}}{\lambda_{\max}}}))$


★ Currently working on A nonsymmetric

Convergence rate. $A \in \mathbb{R}^{400 \times 400}$ symmetric. $f(\lambda) = \lambda^{-1/2}$




$$\sigma(A) = [0.01, 0.9]$$

Uniform spectral distribution



$$\sigma(A) = [1, 50]$$


Experiment. $A \in \mathbb{R}^{400 \times 400}$ normal. $f(\lambda) = \lambda^{-1/2}$



$\sigma(A)$ on an elliptic curve in \mathbb{C}^+ with center on real axis

Log-uniform spectral distribution.

$$f_{(3)}(z) = \exp(-\sqrt{z}), \quad f_{(5)}(z) = z^{-1/2}, \quad f_{(7)}(z) = z^{-1/4}$$




cases *-1: $\lambda_{\min} = 10^{-1}$

cases *-2: $\lambda_{\min} = 10^{-4}$

x-axis: $\lambda_{\max} = 10^k$

SI-Lanczos and dependence on parameter γ



Log-uniform spectral distribution

Comparisons: CPU Time in Matlab

$A \in \mathbb{R}^{4900 \times 4900}$: $\mathcal{L}(u) = -\frac{1}{10}u_{xx} - 100u_{yy}$, in $[0, 1]^2$, hom.b.c.


$\sigma(A) \in [9.6 \cdot 10^2, 1.96 \cdot 10^6]$

Method	space dim.	CPU Time
Standard Krylov	185	16.02
SI-Lanczos(0.001)	62	1.00
SI-Lanczos (1e-5)	49	0.60
SI-Lanczos ($\gamma=2e-5$)	33	0.32
Extended Krylov	32	0.20

No reorthogonalization.

A nonsymmetric matrix. $f(z) = \sqrt{z}$

$A \in \mathbb{R}^{900 \times 900}$: $\mathcal{L}(u) = -100u_{xx} - u_{yy} + 10xu_x$ in $[0, 1]^2$, hom.b.c.



$\sigma(A) \subset \mathbb{R}$, $\lambda_{\min} = 9.2 \cdot 10^2$, $\lambda_{\max} = 3.6 \cdot 10^5$

Full orthogonalization

Conclusions and work in progress

- New implementation of the Extended Krylov method
- Improved convergence bounds for A symmetric
- New convergence results for A nonsymmetric (to be completed)
- Performance:
 - Competitive with respect to available methods
 - Does not depend on parameters
 - Competitive for A nonsymmetric (preliminary tests)