

The Topology of Tiling Spaces

Jean BELLISSARD^{1 2}

Georgia Institute of Technology, Atlanta,
<http://www.math.gatech.edu/~jeanbel/>

Collaborations:

R. BENEDETTI (U. Pisa, Italy)

J.-M. GAMBAUDO (U. Nice, France)

D.J.L. HERRMANN (U. Tübingen, Germany)

J. KELLENDONK (U. Lyon I, Lyon, France)

J. SAVINIEN (Gatech, Atlanta, GA & U. Bielefeld, Germany)

M. ZARROUATI (U. Toulouse III, Toulouse, France)

¹Georgia Institute of Technology, School of Mathematics, Atlanta, GA 30332-0160

²e-mail: jeanbel@math.gatech.edu

Main References

J. BELLISSARD, *The Gap Labeling Theorems for Schrödinger's Operators*,
in *From Number Theory to Physics*, pp. 538-630, Les Houches March 89, Springer,
J.M. Luck, P. Moussa & M. Waldschmidt Eds., (1993).

J. KELLENDONK,
The local structure of tilings and their integer group of coinvariants,
Comm. Math. Phys., **187**, (1997), 1823-1842.

J. E. ANDERSON, I. PUTNAM,
Topological invariants for substitution tilings and their associated C^ -algebras*,
Ergodic Theory Dynam. Systems, **18**, (1998), 509-537.

A. H. FORREST, J. HUNTON,
The cohomology and K-theory of commuting homeomorphisms of the Cantor set,
Ergodic Theory Dynam. Systems, **19**, (1999), 611-625.

J. C. LAGARIAS,
Geometric models for quasicrystals I & II,
Discrete Comput. Geom., **21**, (1999), 161-191 & 345-372.

J. BELLISSARD, D. HERRMANN, M. ZARROUATI,
Hull of Aperiodic Solids and Gap Labeling Theorems,
In *Directions in Mathematical Quasicrystals*, CRM Monograph Series,
Volume 13, (2000), 207-259, M.B. Baake & R.V. Moody Eds., AMS Providence.

L. SADUN, R. F. WILLIAMS,
Tiling spaces are Cantor set fiber bundles,
Ergodic Theory Dynam. Systems, **23**, (2003), 307-316.


J. BELLISSARD, R. BENEDETTI, J. M. GAMBAUDO,
Spaces of Tilings, Finite Telescopic Approximations,
Comm. Math. Phys., **261**, (2006), 1-41.

J. BELLISSARD, J. SAVINIEN,
A Spectral Sequence for the K-theory of Tiling Spaces,
arXiv:0705.2483, submitted to Ergod. Th. Dyn. Syst., 2007.


Content

1. Tilings, Tilings...
2. The Hull as a Dynamical System
3. Branched Oriented Flat Riemannian Manifolds
4. Cohomology and K -Theory
5. Conclusion


I - Tilings, Tilings,...




- A triangle tiling -




- Dominos on a triangular lattice -




- Building the chair tiling -




- The chair tiling -




- The Penrose tiling -




- Kites and Darts -




- Rhombi in Penrose's tiling -




- The Penrose tiling -




- The octagonal tiling -




- Octagonal tiling: inflation rules -




- Another octagonal tiling -



- Another octagonal tiling -



- Building the Pinwheel Tiling -



- The Pinwheel Tiling -

Aperiodic Materials

1. *Periodic Crystals* in d -dimensions:
translation and crystal symmetries.
Translation group $\mathcal{T} \simeq \mathbb{Z}^d$.
2. *Periodic Crystals in a Uniform Magnetic Field*;
magnetic oscillations, Shubnikov-de Haas, de Haas-van Alfen.
The magnetic field breaks the translation invariance to give some quasiperiodicity.

3. *Quasicrystals*: no translation symmetry, but icosahedral symmetry. Ex.:

(a) **Al_{62.5}Cu₂₅Fe_{12.5}**;

(b) **Al₇₀Pd₂₂Mn₈**;


(c) **Al₇₀Pd₂₂Re₈**;

4. *Disordered Media*: random atomic positions


(a) Normal metals (with defects or impurities);

(b) Alloys

(c) Doped semiconductors (**Si**, **AsGa**, ...);



- The icosahedral quasicrystal $AlPdMn$ -



- The icosahedral quasicrystal HoMgZn -

II - The Hull as a Dynamical System

Point Sets

A subset $\mathcal{L} \subset \mathbb{R}^d$ may be:

1. *Discrete*.
2. *Uniformly discrete*: $\exists r > 0$ s.t. each ball of radius r contains at most one point of \mathcal{L} .
3. *Relatively dense*: $\exists R > 0$ s.t. each ball of radius R contains at least one points of \mathcal{L} .
4. A *Delone* set: \mathcal{L} is uniformly discrete and relatively dense.
5. *Finite Local Complexity (FLC)*: $\mathcal{L} - \mathcal{L}$ is discrete and closed.
6. *Meyer* set: \mathcal{L} and $\mathcal{L} - \mathcal{L}$ are Delone.

Point Sets and Point Measures

$\mathfrak{M}(\mathbb{R}^d)$ is the set of Radon measures on \mathbb{R}^d namely the dual space to $C_c(\mathbb{R}^d)$ (continuous functions with compact support), endowed with the weak* topology.

For \mathcal{L} a *uniformly discrete* point set in \mathbb{R}^d :

$$\nu := \nu^{\mathcal{L}} = \sum_{y \in \mathcal{L}} \delta(x - y) \quad \in \mathfrak{M}(\mathbb{R}^d) .$$


Point Sets and Tilings

Given a tiling with finitely many tiles (*modulo translations*), a Delone set is obtained by defining a point in the interior of each (*translation equivalence class of*) tile.


Conversely, given a Delone set, a tiling is built through the *Voronoi cells*

$$V(x) = \{a \in \mathbb{R}^d; |a - x| < |a - y|, \forall y \in \mathcal{L} \setminus \{x\}\}$$

1. $V(x)$ is an *open convex polyhedron* containing $B(x; r)$ and contained into $\overline{B(x; R)}$.
2. Two Voronoi cells touch face-to-face.
3. If \mathcal{L} is *FLC*, then the Voronoi tiling has finitely many tiles modulo translations.



- Building a Voronoi cell-



- A Delone set and its Voronoi Tiling-

The Hull

A point measure is $\mu \in \mathfrak{M}(\mathbb{R}^d)$ such that $\mu(B) \in \mathbb{N}$ for any ball $B \subset \mathbb{R}^d$. Its support is

1. *Discrete*.
2. *r-Uniformly discrete*: iff $\forall B$ ball of radius r , $\mu(B) \leq 1$.
3. *R-Relatively dense*: iff for each ball B of radius R , $\mu(B) \geq 1$.

\mathbb{R}^d acts on $\mathfrak{M}(\mathbb{R}^d)$ by translation.

Theorem 1 *The set of r -uniformly discrete point measures is compact and \mathbb{R}^d -invariant.*

Its subset of R -relatively dense measures is compact and \mathbb{R}^d -invariant.

Definition 1 *Given \mathcal{L} a uniformly discrete subset of \mathbb{R}^d , the Hull of \mathcal{L} is the closure in $\mathfrak{M}(\mathbb{R}^d)$ of the \mathbb{R}^d -orbit of $\nu_{\mathcal{L}}$.*

Hence the Hull is a *compact metrizable space* on which \mathbb{R}^d acts by *homeomorphisms*.

Properties of the Hull

If $\mathcal{L} \subset \mathbb{R}^d$ is r -uniformly discrete with Hull Ω then using compactness

1. each point $\omega \in \Omega$ is an r -uniformly discrete point measure with support \mathcal{L}_ω .
2. if \mathcal{L} is (r, R) -Delone, so are all \mathcal{L}_ω 's.
3. if, in addition, \mathcal{L} is FLC, so are all the \mathcal{L}_ω 's.

Moreover then $\mathcal{L} - \mathcal{L} = \mathcal{L}_\omega - \mathcal{L}_\omega \forall \omega \in \Omega$.

Definition 2 *The transversal of the Hull Ω of a uniformly discrete set is the set of $\omega \in \Omega$ such that $0 \in \mathcal{L}_\omega$.*

Theorem 2 *If \mathcal{L} is FLC, then its transversal is completely discontinuous.*

Local Isomorphism Classes and Tiling Space

A *patch* is a finite subset of \mathcal{L} of the form

$$p = (\mathcal{L} - x) \cap \overline{B(0, r_1)} \quad x \in \mathcal{L}, \quad r_1 \geq 0$$

Given \mathcal{L} a repetitive, FLC, Delone set let \mathcal{W} be its set of finite patches: it is called the *the \mathcal{L} -dictionary*.

A Delone set (or a Tiling) \mathcal{L}' is *locally isomorphic* to \mathcal{L} if it has the same dictionary. The *Tiling Space* of \mathcal{L} is the set of *Local Isomorphism Classes* of \mathcal{L} .

Theorem 3 *The Tiling Space of \mathcal{L} coincides with its Hull.*

Minimality

\mathcal{L} is *repetitive* if for any finite patch p there is $R > 0$ such that each ball of radius R contains an ϵ -approximant of a translated of p .

Theorem 4 \mathbb{R}^d acts minimally on Ω if and only if \mathcal{L} is repetitive.

Examples


1. *Crystals* : $\Omega = \mathbb{R}^d / \mathcal{T} \simeq \mathbb{T}^d$ with the quotient action of \mathbb{R}^d on itself. (Here \mathcal{T} is the translation group leaving the lattice invariant. \mathcal{T} is isomorphic to \mathbb{Z}^D .)

The transversal is a finite set (number of point per unit cell).

2. *Impurities in Si* : let \mathcal{L} be the lattices sites for *Si* atoms (it is a Bravais lattice). Let \mathfrak{A} be a finite set (alphabet) indexing the types of impurities.

The transversal is $X = \mathfrak{A}^{\mathbb{Z}^d}$ with \mathbb{Z}^d -action given by shifts.

The Hull Ω is the mapping torus of X .



- The Hull of a Periodic Lattice -

Quasicrystals

Use the *cut-and-project* construction:


$$\mathbb{R}^d \simeq \mathcal{E}_{\parallel} \xleftarrow{\pi_{\parallel}} \mathbb{R}^n \xrightarrow{\pi_{\perp}} \mathcal{E}_{\perp} \simeq \mathbb{R}^{n-d}$$

$$\mathcal{L} \xleftarrow{\pi_{\parallel}} \tilde{\mathcal{L}} \xrightarrow{\pi_{\perp}} W ,$$


Here

1. $\tilde{\mathcal{L}}$ is a *lattice* in \mathbb{R}^n ,
2. the *window* W is a compact polytope.
3. \mathcal{L} is the *quasilattice* in \mathcal{E}_{\parallel} defined as

$$\mathcal{L} = \{ \pi_{\parallel}(m) \in \mathcal{E}_{\parallel} ; m \in \tilde{\mathcal{L}}, \pi_{\perp}(m) \in W \}$$



– The cut-and-project construction –



- The transversal of the Octagonal Tiling is completely disconnected -

III - Branched Oriented Flat Riemannian Manifolds

Laminations and Boxes

A *lamination* is a foliated manifold with C^∞ -structure along the leaves but only finite C^0 -structure transversally. The *Hull of a Delone set is a lamination* with \mathbb{R}^d -orbits as leaves.

If \mathcal{L} is a *FLC, repetitive, Delone* set, with Hull Ω a *box* is the homeomorphic image of

$$\phi : (\omega, x) \in F \times U \mapsto \tau^{-x}\omega \in \Omega$$

if F is a clopen subset of the transversal, $U \subset \mathbb{R}^d$ is open and τ denotes the \mathbb{R}^d -action on Ω .

A *quasi-partition* is a family $(B_i)_{i=1}^n$ of boxes such that $\bigcup_i \overline{B_i} = \Omega$ and $B_i \cap B_j = \emptyset$.


Theorem 5 *The Hull of a FLC, repetitive, Delone set admits a finite quasi-partition. It is always possible to choose these boxes as $\phi(F \times U)$ with U a d -rectangle.*

Branched Oriented Flat Manifolds


Flattening a box decomposition along the transverse direction leads to a *Branched Oriented Flat manifold*. Such manifolds can be built from the tiling itself as follows

Step 1:

1. X is the disjoint union of all *prototiles*;
2. glue prototiles T_1 and T_2 along a face $F_1 \subset T_1$ and $F_2 \subset T_2$ if F_2 is a translated of F_1 and if there are $x_1, x_2 \in \mathbb{R}^d$ such that $x_i + T_i$ are tiles of \mathcal{T} with $(x_1 + T_1) \cap (x_2 + T_2) = x_1 + F_1 = x_2 + F_2$;
3. after identification of faces, X becomes a *branched oriented flat manifold* (BOF) B_0 .




- Branching -



- Vertex branching for the octagonal tiling -

Step 2:

1. Having defined the patch p_n for $n \geq 0$, let \mathcal{L}_n be the subset of \mathcal{L} of points centered at a translated of p_n . By repetitivity this is a FLC repetitive Delone set too. Its prototiles are tiled by tiles of \mathcal{L} and define a finite family \mathfrak{P}_n of patches.
2. Each patch in $\mathcal{T} \in \mathfrak{P}_n$ will be collared by the patches of \mathfrak{P}_{n-1} touching it from outside along its frontier. Call such a patch *modulo translation* a *collared patch* and \mathfrak{P}_n^c their set.
3. Proceed then as in Step 1 by replacing prototiles by collared patches to get the BOF-manifold B_n .
4. Then choose p_{n+1} to be the collared patch in \mathfrak{P}_n^c containing p_n .



- A collared patch -

Step 3:

1. Define a *BOF-submersion* $f_n : B_{n+1} \mapsto B_n$ by identifying patches of order n in B_{n+1} with the prototiles of B_n . Note that $Df_n = \mathbf{1}$.
2. Call Ω the *projective limit* of the sequence

$$\dots \xrightarrow{f_{n+1}} B_{n+1} \xrightarrow{f_n} B_n \xrightarrow{f_{n-1}} \dots$$

3. X_1, \dots, X_d are the commuting constant vector fields on B_n generating local translations and giving rise to a \mathbb{R}^d action τ on Ω .

Theorem 6 *The dynamical system*

$$(\Omega, \mathbb{R}^d, \tau) = \varprojlim (B_n, f_n)$$

obtained as inverse limit of branched oriented flat manifolds, is conjugate to the Hull of the Delone set of the tiling \mathcal{T} by an homeomorphism.

IV - Cohomology and K -Theory

Čech Cohomology of the Hull

Let \mathcal{U} be an *open covering* of the Hull. If $U \in \mathcal{U}$, $\mathcal{F}(U)$ is the space of integer valued locally constant function on U .

For $n \in \mathbb{N}$, the n -chains are the element of $C^n(\mathcal{U})$, namely the *free abelian group* generated by the elements of $\mathcal{F}(U_0 \cap \cdots \cap U_n)$ when the U_i varies in \mathcal{U} . A differential is defined by

$$d : C^n(\mathcal{U}) \mapsto C^{n+1}(\mathcal{U})$$

$$df\left(\bigcap_{i=0}^{n+1} U_i\right) = \sum_{j=0}^n (-1)^j f\left(\bigcap_{i:i \neq j} U_i\right)$$

This defines a *complex* with cohomology $\check{H}^n(\mathcal{U}, \mathbb{Z})$. The Čech cohomology group of the Hull Ω is defined as

$$\check{H}^n(\Omega, \mathbb{Z}) = \varinjlim_{\mathcal{U}} \check{H}^n(\mathcal{U}, \mathbb{Z})$$

with ordering given by *refinement* on the set of open covers.

Longitudinal (co)-Homology

J. BELLISSARD, R. BENEDETTI, J.-. GAMBAUDO, Commun. Math. Phys., **261**, (2006), 1-41.

J. KAMINKER, I. PUTNAM, Michigan Math. J., **51**, (2003), 537-546.

M. BENAMEUR, H. OYONO-OYONO, C. R. Math. Acad. Sci. Paris, **334**, (2002), 667-670.

The Homology groups are defined by the inverse limit

$$H_*(\Omega, \mathbb{R}^d) = \varprojlim (H_*(B_n, \mathbb{R}), f_n^*)$$

Theorem 7 (JB, Benedetti, Gambaudo) *The homology group $H_d(\Omega, \mathbb{R}^d)$ admits a canonical positive cone induced by the orientation of \mathbb{R}^d , isomorphic to the affine set of positive \mathbb{R}^d -invariant measures on Ω .*

The cohomology groups are defined by the direct limit

$$H^*(\Omega, \mathbb{R}^d) = \varinjlim (H^*(B_n, \mathbb{R}), f_n^*)$$

The following result is known as the *Gap labeling Theorem* and was proved simultaneously by KAMINKER-PUTNAM, BENAMEUR & OYONO-OYONO, JB-BENDETTI-GAMBAUDO. It is an extension of the *Connes index theorem* for foliations

Theorem 8 *If \mathbb{P} is an \mathbb{R}^d -invariant probability on Ω , then the pairing with $H^d(\Omega, \mathbb{R}^d)$ satisfies*

$$\langle \mathbb{P} | H^d(\Omega, \mathbb{R}^d) \rangle = \int_{\Xi} d\mathbb{P}_{\text{tr}} \, C(\Xi, \mathbb{Z})$$

where Ξ is the transversal, \mathbb{P}_{tr} is the probability on Ξ induced by \mathbb{P} and $C(\Xi, \mathbb{Z})$ is the space of integer valued continuous functions on Ξ .

Pattern-Equivariant Cohomology

J. KELLENDONK, J. Phys. **A36**, (2003), 5765-5772.

J. KELLENDONK, I. PUTNAM, Math. Ann. **334**, (2006), 693-711.

L. SADUN, *Pattern-Equivariant Cohomology with Integer Coefficients* (2007)

Let \mathcal{L} be an FLC, repetitive Delone set in \mathbb{R}^d . A function $f : \mathbb{R}^d \mapsto X$ is *\mathcal{L} -pattern-equivariant* if there is $r > 0$ such that $f(x) = f(y)$ whenever $B(0; r) \cap (\mathcal{L} - x) = B(0; r) \cap (\mathcal{L} - y)$.

The Voronoi tiling of \mathcal{L} can be seen as a *chain complex*, with tiles being the d -cells, and their k -faces being the k -cells.

A *k-cochain* with integer coefficients is then a linear map α defined on the free abelian group of k -chains with values in \mathbb{Z} .

Let $C_{\mathcal{P}}^k(\mathcal{L})$ be the abelian group of \mathcal{L} -pattern equivariant k -cochains. The usual coboundary operator (*de Rham differential*)

$$d_n : C_{\mathcal{P}}^n(\mathcal{L}) \mapsto C_{\mathcal{P}}^{n+1}(\mathcal{L})$$

defines the *\mathcal{L} -pattern equivariant cohomology* denoted by

$$H_{\mathcal{P}}^k(\mathcal{L}, \mathbb{Z}) = \text{Ker } d_n / \text{Im } d_{n-1}$$

The PV-Cohomology

J. BELLISSARD, J.SAVINIEN, *arXiv*: 0705.2483, (2007).

Each cell of the *Voronoi complex* is punctured. The set \mathcal{L}_s of such punctures defines the *simplicial transversal* Ξ_s . An equivalent class, modulo translation, of n -cell σ defines a compact subset $\Xi_s(\sigma)$. χ_σ denotes the characteristic function of $\Xi_s(\sigma)$.

If σ is such a cell and τ belongs to its boundary, then there is a unique vector $x_{\sigma\tau}$ joining the puncture of τ to the one of σ . Correspondingly the translation $T^{x_{\sigma\tau}}$ in the Hull sends $\Xi_s(\tau)$ into a part of $\Xi_s(\sigma)$, defining the translation operator

$$\theta_{\sigma\tau} = \chi_\sigma T^{x_{\sigma\tau}} \chi_\tau$$

where χ_σ denotes the characteristic function of $\Xi_s(\sigma)$.

A *PV- n -cochain* will be a group homomorphism from the group of (oriented) n -chains on the BOF manifold B_0 into the group $C(\Xi_s, \mathbb{Z})$. The Pimsner differential is defined by

$$df(\sigma) = \sum_{\tau \in \partial\sigma} [\sigma : \tau] f(\tau) \circ \theta_{\sigma\tau}$$

Here $[\sigma : \tau]$ denoted the *incidence number* of τ relative to σ . The associate cohomology is $H_P^n(B_0, C(\Xi_s, \mathbb{Z}))$.

Cohomology and K-theory

The main topological property of the Hull (or tiling space) is summarized in the following

Theorem 9 (i) *The various cohomologies, Čech, longitudinal, pattern-equivariant and PV, are isomorphic.*

(ii) *There is a spectral sequence converging to the K-group of the Hull with page 2 given by the cohomology of the Hull.*

(iii) *In dimension $d \leq 3$ the K-group coincides with the cohomology.*

Conclusion

1. *Tilings* can be equivalently be represented by *Delone sets* or *point measures*.
2. The *Hull* allows to give tilings the structure of a *dynamical system* with a transversal.
3. This dynamical system can be seen as a *lamination* or, equivalently, as the *inverse limit* of *Branched Oriented Flat Riemannian Manifolds*.

4. The Čech cohomology is equivalent to the longitudinal one, obtained by inverse limit, to the pattern-equivariant one or to the Pimsner cohomology are equivalent *Cohomology* of the Hull. The *K-group* of the Hull can be computed through a spectral sequence with the cohomology in page 2.
5. In maximum degree, the *Homology* gives the family of *invariant measures* and the *Gap Labelling Theorem*.