

THE DIRICHLET SPACE: A SURVEY

NICOLA ARCOZZI, RICHARD ROCHBERG, ERIC T. SAWYER, AND BRETT D. WICK

ABSTRACT. In this paper we survey many results on the Dirichlet space of analytic functions. Our focus is more on the classical Dirichlet space on the disc and not the potential generalizations to other domains or several variables. Additionally, we focus mainly on certain function theoretic properties of the Dirichlet space and omit covering the interesting connections between this space and operator theory. The results discussed in this survey show what is known about the Dirichlet space and compares it with the related results for the Hardy space.

CONTENTS

1. Introduction	2
2. The Dirichlet Space	2
2.1. The Definition	2
2.2. The Definition in terms of Boundary Values and other Characterizations of the Dirichlet Norm	5
2.3. The Reproducing Kernel	6
3. Carleson measures	6
3.1. Definition and the Capacitary Characterization	6
3.2. Characterizations by Testing Conditions	9
4. The Tree Model	12
4.1. The Bergman Tree	12
4.2. Detour: The Boundary of the Tree and its Relation with the Disc's Boundary.	13
4.3. A Version of the Dirichlet Space on the Tree	14
4.4. Carleson Measures and All That, on the Tree and on the Disc	15
5. The Complete Nevanlinna-Pick property	18
6. The Multiplier Space and other Spaces intrinsic to \mathcal{D} Theory	21
6.1. Multipliers	21
6.2. The Weakly Factored Space $\mathcal{D} \odot \mathcal{D}$ and its Dual	22
6.3. The Corona Theorem	23
7. Interpolating sequences	26
7.1. Interpolating Sequences for \mathcal{D} and its Multiplier Space	27
7.2. Weak Interpolation and "Onto" Interpolation	29
References	30

N.A.'s work partially supported by the COFIN project Analisi Armonica, funded by the Italian Minister for Research.

R.R.'s work supported by the National Science Foundation under Grant No. 0700238.

E.S.'s work supported by the National Science and Engineering Council of Canada.

B.W.'s work supported by the National Science Foundation under Grant No. 1001098.

1. INTRODUCTION

Notation. The unit disc will be denoted by $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$ and the unit circle by $\mathbb{S} = \partial\mathbb{D}$. If Ω is open in \mathbb{C} , $H(\Omega)$ is the space of the functions which are holomorphic in Ω . A function $\varphi : \mathbb{S} \rightarrow \mathbb{C}$ is identified with a function defined on $[0, 2\pi)$; $\varphi(e^{i\theta}) = \varphi(\theta)$.

Given two (variable) quantities A and B , we write $A \approx B$ if there are universal constants $C_1, C_2 > 0$ such that $C_1 A \leq B \leq C_2 A$. Similarly, we use the symbol \lesssim . If A_1, \dots, A_n are mathematical objects, the symbol $C(A_1, \dots, A_n)$ denotes a constant which only depends on A_1, \dots, A_n .

The Dirichlet space, together with the Hardy and the Bergman space, is one of the three classical spaces of holomorphic functions in the unit disc. Its theory is old, but over the past thirty years much has been learned about it and about the operators acting on it. The aim of this article is to survey some aspects, old and new, of the Dirichlet theory.

We will concentrate on the “classical” Dirichlet space and we will not dwell into its interesting extensions and generalizations. The only exception, because instrumental to our discourse, will be some discrete function spaces on trees.

Our main focus will be a *Carleson-type* program which has been unfolding over the past thirty years. To have, that is, a knowledge of Dirichlet space comparable to that we have, mainly due to the work of Lennart Carleson, for the Hardy space H^2 : weighted imbedding theorems (“Carleson measures”); interpolating sequences; the Corona Theorem. We also consider other topics which are well understood in the Hardy case: boundary behavior; bilinear forms; applications of Nevanlinna-Pick theory; spaces which are necessary to develop the Hilbert space theory (H^1 and BMO , for instance, in the case of H^2). Let us further mention a topic which is specifically related to the Dirichlet theory: the rich relationship with potential theory and capacity.

This survey is much less than comprehensive. We will be mainly focused on the properties of the Dirichlet space *per se*, and we will talk about the rich operator theory which has been developing on it during the past twenty years or so, only when this intersects our main path. We are also biased, more or less voluntarily, towards the topics on which we have been working, with more or less success. If the scope of the survey is narrow, we will try to give some detail of the ideas and arguments, in the hope to render a service to those who for the first time approach the subject.

Let us finally mention the excellent survey on the Dirichlet space [44], to which we direct the reader for the discussion on the local Dirichlet integral, Carleson’s and Douglas’ formulas and the discussion of the invariant subspaces. Also, [44] contains a discussion of zero sets and boundary behavior, hence we will only tangentially touch on these topics here. The article [59] surveys some results in the operator theory on the Dirichlet space.

2. THE DIRICHLET SPACE

2.1. The Definition. The *Dirichlet space* \mathcal{D} is the Hilbert space of the functions f analytic in the unit disc $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$ for which the semi-norm

$$(1) \quad \|f\|_{\mathcal{D},*}^2 = \int_{\mathbb{D}} |f'(z)|^2 dA(z)$$

is finite. Here, $dA(x+iy) = \frac{1}{\pi} dx dy$ is normalized area measure. An easy calculation with Fourier coefficients shows that, if $f(z) = \sum_{n=0}^{\infty} a_n z^n$,

$$(2) \quad \|f\|_{\mathcal{D},*}^2 = \sum_{n=1}^{\infty} n |a_n|^2.$$

The Dirichlet space sits then inside the analytic Hardy space H^2 . In particular, Dirichlet functions have nontangential limits at *a.e.* point on the boundary of \mathbb{D} . We will see later that much more can be said, both on the kind of approach region and on the size of the exceptional set.

There are different ways to make the semi-norm into a norm. Here, we use as norm and inner product, respectively,

$$(3) \quad \begin{aligned} \|f\|_{\mathcal{D}}^2 &= \|f\|_{\mathcal{D},*}^2 + \|f\|_{H^2(\mathbb{S})}^2, \\ \langle f, g \rangle_{\mathcal{D}} &= \langle f, g \rangle_{\mathcal{D},*} + \langle f, g \rangle_{H^2(\mathbb{S})} \\ &= \int_{\mathbb{D}} f'(z) \overline{g'(z)} dA(z) + \frac{1}{2\pi} \int_0^{2\pi} f(e^{i\theta}) \overline{g(e^{i\theta})} d\theta. \end{aligned}$$

Another possibility is to let $\|f\|_{\mathcal{D}}^2 = \|f\|_{\mathcal{D},*}^2 + |f(0)|^2$. Most analysis on \mathcal{D} carries out in the same way, no matter what is the chosen norm. There is an important exception to this rule. The *Complete Nevanlinna-Pick Property* is not invariant under change of norm: it is satisfied by $\|\cdot\|_{\mathcal{D}}$, not by $\|\cdot\|_{\mathcal{D},*}$.

The Dirichlet semi-norm has two different, interesting geometric interpretations.

(E) Since $Jf = |f'|^2$ is the Jacobian determinant of f ,

$$(4) \quad \|f\|_{\mathcal{D},*}^2 = \int_{\mathbb{D}} dA(f(z)) = A(f(\mathbb{D}))$$

is the area of the image of f , counting multiplicities. This invariance property, which depends on the *values* of functions in \mathcal{D} , implies that the Dirichlet class is invariant under biholomorphisms of the disc.

(H) Let $ds^2 = \frac{|dz|^2}{(1-|z|^2)^2}$ be the hyperbolic metric in the unit disc. The (normalized) hyperbolic area density is $d\lambda(z) = \frac{dA(z)}{(1-|z|^2)^2}$ and the intrinsic derivative of a holomorphic $f : (\mathbb{D}, ds^2) \rightarrow (\mathbb{C}, |dz|^2)$ is $\delta f(z) = (1-|z|^2)|f'(z)|$. Then,

$$(5) \quad \|f\|_{\mathcal{D},*}^2 = \int_{\mathbb{D}} (\delta f)^2 d\lambda$$

is defined in purely hyperbolic terms.

Since any Blaschke product with n factors in an n -to-1 covering of the unit disc, (E) implies that the Dirichlet space only contains finite Blaschke products. The characterization of the zero sets for the Dirichlet space is a difficult, open problem. On the positive side, (E) allows one to define the Dirichlet space on any simply connected domain $\Omega \subsetneq \mathbb{C}$,

$$\|f\|_{\mathcal{D}(\Omega),*}^2 := \int_{\Omega} |f'(z)|^2 dA(z) = \|f \circ \varphi\|_{\mathcal{D},*}^2,$$

where φ is any conformal map of the unit disc onto Ω . In particular, this shows that the Dirichlet semi-norm is invariant under the Möbvius group $\mathcal{M}(\mathbb{D})$.

Infinite Blaschke products provide examples of bounded functions which are not in the Dirichlet space. On the other hand, conformal maps of the unit disc

onto unbounded regions having finite area provide examples of unbounded Dirichlet functions.

The group $\mathcal{M}(\mathbb{D})$ acts on (\mathbb{D}, ds^2) as the group of the sense preserving isometries. It follows from (H) as well, then, the conformal invariance of the Dirichlet semi-norm: $\|f \circ \varphi\|_{\mathcal{D},*} = \|f\|_{\mathcal{D},*}$ when $\varphi \in \mathcal{M}(\mathbb{D})$. In fact, Arazy and Fischer [6] showed that the Dirichlet semi-norm is the only Möbius invariant, Hilbert semi-norm for functions holomorphic in the unit disc. Also, the Dirichlet space is the only Möbius invariant Hilbert space of holomorphic functions on the unit disc. Sometimes it is preferable to use the *pseudo-hyperbolic* metric instead,

$$\rho(z, w) := \left| \frac{z - w}{1 - \bar{w}z} \right|.$$

The hyperbolic metric d and the pseudo-hyperbolic metric are functionally related,

$$d = \frac{1}{2} \log \frac{1 + \rho}{1 - \rho}, \quad \rho = \frac{e^d - e^{-d}}{e^d + e^{-d}}.$$

(The hyperbolic metric is the only Riemannian metric which coincides with the pseudo-hyperbolic metric when infinitesimally small). The triangle property for the hyperbolic metric is equivalent to an enhanced triangle property for the pseudo-hyperbolic metric:

$$\rho(z, w) \leq \frac{\rho(z, t) + \rho(t, w)}{1 + \rho(z, t)\rho(t, w)}.$$

We conclude with a simple and entertaining consequence of (H). The isoperimetric inequality

$$(6) \quad Area(\Omega) \leq \frac{1}{4\pi} [Length(\partial\Omega)]^2$$

is equivalent, by Riemann's Mapping Theorem and by the extension of (6) itself to areas with multiplicities, to the inequality

$$\|f\|_{\mathcal{D},*}^2 = \int_{\mathbb{D}} |f'|^2 dA \leq \left[\frac{1}{2\pi} \int_{\partial\mathbb{D}} |f'(e^{i\theta})| d\theta \right]^2 = \|f'\|_{H^1}^2.$$

Set $f' = g$ in the last inequality: the isoperimetric inequality becomes the imbedding of the Bergman space A^2 in H^1 , with optimal constant:

$$\|g\|_{A^2}^2 \leq \|g\|_{H^1}^2,$$

the constant functions being extremal.

2.1.1. *The Hardy case space H^2 .* The “classical” Hilbert spaces of holomorphic functions on the unit disc are the Dirichlet space just introduced, the Bergman space A^2 ,

$$\|f\|_{A^2}^2 = \int_{\mathbb{D}} |f(z)|^2 dA(z),$$

and the Hardy space H^2 ,

$$\|f\|_{H^2}^2 = \sup_{0 < r < 1} \frac{1}{2\pi} \int_0^{2\pi} |f(re^{i\theta})|^2 d\theta.$$

The Hardy space is especially important because of its direct rôle in operator theory, as a prototype for the study of boundary problems for elliptic differential equations, for its analogy with important probabilistic objects (martingales), and for many other reasons. It has been studied in depth and its theory has become a model

for the theory of other classical and not so classical function spaces. Many results surveyed in this article have been first proved, in a different version, for the Hardy space.

It is interesting to observe that both the Hardy and the Bergman space can be thought of as weighted Dirichlet spaces. We consider here the case of the Hardy space. If $f(0) = 0$, then

$$\|f\|_{H^2}^2 = \int_{\mathbb{D}} |f'(z)|^2 \log(|z|^{-2}) dA(z) \approx \int_{\mathbb{D}} |f'(z)|^2 (1 - |z|^2) dA(z).$$

This representation of the Hardy functions is more than a curiosity. Since H^2 is a reproducing kernel Hilbert space (RKHS) of functions, we are interested in having a norm which depends on the values of f in the *interior* of the unit disc. (Indeed, the usual norm is in terms of interior values as well, although through the mediation of sup).

2.2. The Definition in terms of Boundary Values and other Characterizations of the Dirichlet Norm. Let $\mathbb{S} = \partial\mathbb{D}$ be the unit circle and $\mathcal{H}^{1/2}(\mathbb{S})$ be the fractional Sobolev space containing the functions $\varphi \in L^2(\mathbb{S})$ having “1/2” derivative in $L^2(\mathbb{S})$. More precisely, if $\varphi(\theta) = \sum_{n=1}^{+\infty} [a_n \cos(n\theta) + b_n \sin(n\theta)]$, then the $\mathcal{H}^{1/2}(\mathbb{S})$ semi-norm of φ is

$$(7) \quad \|\varphi\|_{\mathcal{H}^{1/2}(\mathbb{S})}^2 = \sum_{n=1}^{+\infty} n(|a_n|^2 + |b_n|^2).$$

By definition,

$$\mathcal{D} \equiv \mathcal{H}^{1/2}(\mathbb{S}) \cap H(\mathbb{D}).$$

This expresses the fact that, restricting Sobolev functions from the plane to smooth curves, “there is a loss of 1/2 derivative”.

2.2.1. The Definition of Rochberg and Wu. In [43], Rochberg and Wu gave a characterization of the Dirichlet norm in terms of difference quotients of the function.

Theorem 1 (Rochberg-Wu, [43]). *Let $\sigma, \tau > -1$. For a function f analytic on the unit disc we have the semi-norm equivalence:*

$$\|f\|_{\mathcal{D},*}^2 \approx \int_{\mathbb{D}} \int_{\mathbb{D}} \frac{|g(z) - g(w)|^2}{|1 - z\bar{w}|^{\sigma+\tau+4}} (1 - |z|^2)^\sigma (1 - |w|^2)^\tau dA(w) dA(z).$$

For $\sigma = \tau = 1/2$, the Theorem holds with equality instead of approximate equality; see [7]. The result in [43] extends to weighted Dirichlet spaces and, with a different, discrete-like proof, to analytic Besov spaces [23]. The characterization in Theorem 1 is similar in spirit to the usual boundary characterization for functions in $\mathcal{H}^{1/2}(\mathbb{S})$:

$$\|\varphi\|_{\mathcal{H}^{1/2}(\mathbb{S})}^2 \approx \int_0^{2\pi} \int_0^{2\pi} \frac{|\varphi(\zeta) - \varphi(\xi)|^2}{|\zeta - \xi|^2} d\zeta d\xi$$

2.2.2. The characterization of Bøe. In [24], Bøe obtained an interesting characterization of the norm in analytic Besov spaces, in terms of the mean oscillation of the function’s modulus, with respect to harmonic measure. We give Bøe’s result in the Dirichlet case.

Theorem 2 (Böe, [24]). *For $z \in \mathbb{D}$, let*

$$d\mu_z(\zeta) = \frac{1}{2\pi} \frac{1 - |z|^2}{|\zeta - z|^2} d\zeta$$

be harmonic measure on \mathbb{S} with respect to z . Then,

$$\|f\|_{\mathcal{D},*}^2 \approx \int_{\mathbb{D}} \left(\int_0^{2\pi} |f(e^{i\zeta})| d\mu_z(\zeta) - |f(z)| \right)^2 \frac{dA(z)}{(1 - |z|^2)^2}.$$

2.3. The Reproducing Kernel. The space \mathcal{D} has bounded point evaluation $\eta_z : f \mapsto f(z)$ at each point $z \in \mathbb{D}$. Equivalently, it has a reproducing kernel. In fact, it is easily checked that

$$f(z) = \langle f, K_z \rangle_{\mathcal{D}}, \text{ with } K_z(w) = \frac{1}{\bar{z}w} \log \frac{1}{1 - \bar{z}w}.$$

(For the norm $\|\cdot\|_{\mathcal{D}}$ introduced earlier, the reproducing kernel is

$$\tilde{K}_z(w) = 1 + \log \frac{1}{1 - \bar{z}w}.$$

which is comfortable in estimates for the integral operator having $\tilde{K}_z(w)$ as kernel).

It is a general fact that $\|\eta_z\|_{\mathcal{D}^*} = \|K_z\|_{\mathcal{D}}$ and an easy calculation gives

$$\|K_z\|_{\mathcal{D}}^2 \approx 1 + \log \frac{1}{1 - |z|} \approx 1 + d(z, 0).$$

More generally, we have that functions in the Dirichlet space are Hölder continuous of order $1/2$ with respect to the hyperbolic distance:

$$(8) \quad |f(z) - f(w)| \leq C \|f\|_{\mathcal{D},*} d(z, w)^{1/2}.$$

The reproducing kernel $K_z(w) = K(z, w)$ satisfies some estimates which are important in applications and reveal its geometric nature.

- (a) $\Re K(z, w) \approx |K(z, w)|$.
- (b) Let $z \wedge w$ be the point which is closest to the origin (in either hyperbolic or Euclidean metric) on the hyperbolic geodesic joining z and w . Then,

$$\Re K(z, w) \approx d(0, z \wedge w) + 1.$$

- (c) $\frac{d}{dw} K(z, w) = \frac{\bar{z}}{1 - \bar{z}w}$, and we have:

- (c1) $\Re \frac{1}{1 - \bar{z}w} \geq 0$ for all z, w in \mathbb{D} ;
- (c2) $\Re \frac{1}{1 - \bar{z}w} \approx (1 - |z|^2)^{-1}$ for $w \in S(z)$, where

$$S(z) = \{w \in \mathbb{D} : |1 - \bar{z}w| \leq 1 - |z|^2\}$$

is the Carleson box with centre z .

3. CARLESON MEASURES

3.1. Definition and the Capacitary Characterization. A positive Borel measure μ on $\bar{\mathbb{D}}$ is called a *Carleson measure* for the Dirichlet space if for some finite $C > 0$

$$(9) \quad \int_{\bar{\mathbb{D}}} |f|^2 d\mu \leq C \|f\|_{\mathcal{D}}^2 \quad \forall f \in \mathcal{D}.$$

The smallest C in (9) is the *Carleson measure norm* of μ and it will be denoted by $[\mu] = [\mu]_{CM(\mathcal{D})}$. The space of the Carleson measures for \mathcal{D} is denoted by $CM(\mathcal{D})$. Carleson measures supported on the boundary could be thought of as substitutes

for point evaluation (which is not well defined at boundary points). By definition, in fact, the function f exists, in a quantitative way, on sets which support a strictly positive Carleson measure. It is then to be expected that there is a relationship between Carleson measures and boundary values of Dirichlet functions. This is explained further below.

Carleson measures proved to be a central concept in the theory of the Dirichlet space in many other ways. Let us mention:

- Multipliers,
- Interpolating Sequences,
- Bilinear Forms,
- Boundary Values.

Since Carleson measures play such an important role, it is important then to have efficient ways to characterize them. The first such characterization was given by Stegenga [50] in terms of capacity.

We first introduce the Riesz-Bessel kernel of order $1/2$ on \mathbb{S} ,

$$(10) \quad k_{\mathbb{S},1/2}(\theta, \eta) = |\theta - \eta|^{-1/2},$$

where the difference $\theta - \eta \in [-\pi, \pi]$ is taken modulo π . The kernel extends to a convolution operator, which we still call $k_{\mathbb{S},1/2}$, acting on Borel measures supported on \mathbb{S} ,

$$k_{\mathbb{S},1/2}\nu(\theta) = \int_{\mathbb{S}} k_{\mathbb{S},1/2}(\theta - \eta) d\nu(\eta).$$

Let $E \subseteq \mathbb{S}$ be a closed set. The $(\mathbb{S}, 1/2)$ -Bessel capacity of E is

$$(11) \quad \text{Cap}_{\mathbb{S},1/2} := \inf \left\{ \|h\|_{L^2(\mathbb{S})}^2 : h \geq 0 \text{ and } k_{1/2,\mathbb{S}}h \geq 1 \text{ on } E \right\}.$$

It is a well know fact [51] that $\|k_{\mathbb{S},1/2}h\|_{\mathcal{H}^{1/2}(\mathbb{S})} \approx \|h\|_{L^2(\mathbb{S})}$, i.e. that $h \mapsto k_{\mathbb{S},1/2}h$ is an approximate isometry of $L^2(\mathbb{S})$ into $\mathcal{H}^{1/2}(\mathbb{S})$. Hence,

$$\text{Cap}_{\mathbb{S},1/2} \approx \inf \left\{ \|\varphi\|_{\mathcal{H}^{1/2}(\mathbb{S})}^2 : (k_{\mathbb{S},1/2})^{-1}\varphi \geq 0 \text{ and } \varphi \geq 1 \text{ on } E \right\}.$$

Theorem 3 (Stegenga, [50]). *Let $\mu \geq 0$ be a positive, Borel measure on \mathbb{D} . Then μ is Carleson for \mathcal{D} if and only if there is a positive constant $C(\mu)$ such that, for any choice of finitely many disjoint, closed arcs $I_1, \dots, I_n \subseteq \mathbb{S}$, we have that*

$$(12) \quad \mu(\cup_{i=1}^n S(I_i)) \leq C(\mu) \text{Cap}_{\mathbb{S},1/2}(\cup_{i=1}^n I_i).$$

Moreover, $C(\mu) \approx [\mu]_{CM(\mathcal{D})}$ estimates the best value of the constant $C(\mu)$.

It is expected that capacity plays a rôle in the theory of the Dirichlet space. In fact, as we have seen, the Dirichlet space is intimately related to at least two Sobolev spaces ($\mathcal{H}^{1/2}(\mathbb{S})$ and $\mathcal{H}^1(\mathbb{C})$, which is defined below), and capacity plays in Sobolev theory the rôle played by Lebesgue measure in the theory of Hardy spaces. In Dirichlet space theory, this fact has been recognized for a long time (see, for instance, [20]); actually, before Sobolev theory reached maturity.

It is a useful exercise comparing Stegenga's capacity condition and Carleson's condition for the Carleson measures for the Hardy space. Carleson [28] proved that, for a positive, Borel measure μ on \mathbb{D} ,

$$\int_{\mathbb{D}} |f|^2 d\mu \leq C(\mu) \|f\|_{H^2}^2 \iff \mu(S(I)) \leq C'(\mu)|I|,$$

for all closed sub-arcs I of the unit circle. Moreover, the best constants in the two inequalities are comparable. In some sense, Carleson's characterization says that μ satisfies the imbedding $H^2 \hookrightarrow L^2(\mu)$ if and only if it behaves (no worse than) the arclength measure on \mathbb{S} , the measure underlying the Hardy theory. We could also "explain" Carleson's condition in terms of the reproducing kernel for the Hardy space,

$$K_z^{H^2}(w) = \frac{1}{1 - \bar{z}w}, \quad \|K_z^{H^2}\|_{H^2}^2 \approx (1 - |z|)^{-1}.$$

Let I_z be the arc having center in $z/|z|$ and arclength $2\pi(1 - |z|)$. Carleson's condition can then be rephrased as

$$\mu(S(I_z)) \leq C(\mu) \|K_z^{H^2}\|_{H^2}^2.$$

Similar conditions hold for the (weighted) Bergman spaces. One might expect that a necessary and sufficient condition for a measure to belong to $CM(\mathcal{D})$ might be

$$(13) \quad \mu(S(I_z)) \leq C(\mu) \|K_z\|_{\mathcal{D}} \approx \frac{1}{\log \frac{2}{1-|z|}} \approx \text{Cap}_{\mathbb{S}, 1/2}(I_z).$$

The "simple condition" (13) is necessary, but not sufficient. Essentially, this fact is related to the fact that the simple condition does not "add up": if I_j , $j = 1, \dots, 2^n$ are adjacent arcs having the same length, and I is their union, then

$$\sum_j \text{Cap}_{\mathbb{S}, 1/2}(I_j) \approx \frac{2^n}{(\log 2)n + \log \frac{4\pi}{|I|}} \gg \log \frac{1}{\frac{4\pi}{|I|}} \approx \text{Cap}_{\mathbb{S}, 1/2}(I).$$

Stegenga's Theorem has counterparts in the theory of Sobolev spaces, where the problem is that of finding necessary and sufficient conditions on a measure μ so that a *trace inequality* holds. For instance, consider the case of the Sobolev space $\mathcal{H}^1(\mathbb{R}^n)$, containing those functions $h : \mathbb{R}^n \rightarrow \mathbb{C}$ with finite norm

$$\|h\|_{\mathcal{H}^1(\mathbb{R}^n)}^2 = \|h\|_{L^2(\mathbb{R}^n)}^2 + \|\nabla h\|_{L^2(\mathbb{R}^n)}^2,$$

the gradient being the distributional one. The positive, Borel measure μ on \mathbb{R}^n satisfies a trace inequality for $\mathcal{H}^1(\mathbb{R}^n)$ if the imbedding inequality

$$(14) \quad \int_{\mathbb{R}^n} |h|^2 d\mu \leq C(\mu) \|h\|_{\mathcal{H}^1(\mathbb{R}^n)}^2$$

holds. It turns out that (14) is equivalent to the condition that

$$(15) \quad \mu(E) \leq C(\mu) \text{Cap}_{\mathcal{H}^1(\mathbb{R}^n)}(E)$$

holds for all compact subsets $E \subseteq \mathbb{R}^n$. Here, $\text{Cap}_{\mathcal{H}^1(\mathbb{R}^n)}(E)$ is the capacity naturally associated with the space $\mathcal{H}^1(\mathbb{R}^n)$.

There is an extensive literature on trace inequalities, which is closely related to the study of Carleson measures for the Dirichlet space and its extensions. We will not discuss it further, but instead direct the interested reader to [38], [1], [34] and [35], just for a first approach to the subject, from different perspective.

Complex analysts are more familiar with the logarithmic capacity, than with Bessel capacities. It is a classical fact that, for subsets E of the unit circle (or for the real line)

$$(16) \quad \text{Cap}_{\mathbb{S}, 1/2}(E) \approx \log \gamma(E)^{-1},$$

where $\gamma(E)$ is the *logarithmic capacity* (the *transfinite diameter*) of the set E .

3.2. Characterizations by Testing Conditions. The capacity condition has to be checked over all finite unions of arcs. It is natural to wonder whether there is a “single box” condition characterizing the Carleson measures. In fact, there is a string of such conditions, which we are now going to discuss. The following statement rephrases the characterization given in [15]. Let $k(z, w) = \Re K(z, w)$.

Theorem 4 (Arcozzi-Rochberg-Sawyer, [15]). *Let μ be a positive, Borel measure on $\overline{\mathbb{D}}$. Then, μ is a Carleson measure for \mathcal{D} if and only if μ is finite and*

$$(17) \quad \int_{\overline{S(\zeta)}} \int_{\overline{S(\zeta)}} k(z, w) d\mu(w) d\mu(z) \leq C(\mu) \mu(\overline{S(\zeta)})$$

for all ζ in \mathbb{D} .

Moreover, if $C_{\text{best}}(\mu)$ is the best constant in (17), then

$$[\mu]_{CM(\mu)} \approx C_{\text{best}}(\mu) + \mu(\overline{\mathbb{D}}).$$

The actual result in [15] is stated differently. There, it is shown that $\mu \in CM(\mathcal{D})$ if and only if μ is finite and

$$(18) \quad \int_{S(\zeta)} \mu(\overline{S(z)} \cap \overline{S(\zeta)})^2 \frac{dA(z)}{(1 - |z|^2)^2} \leq C(\mu) \mu(\overline{S(\zeta)}),$$

with $[\mu]_{CM(\mu)} \approx C_{\text{best}}(\mu) + \mu(\overline{\mathbb{D}})$. The equivalence between these two conditions will be discussed below, when we will have at our disposal the simple language of trees.

Proof discussion. The basic tools are a duality argument and two weight inequalities for positive kernels. It is instructive to enter in some detail the duality arguments. The definition of Carleson measure says that the imbedding

$$Id : \mathcal{D} \hookrightarrow L^2(\mu)$$

is bounded. Passing to the adjoint $\Theta = Id^*$, this is equivalent to the boundedness of

$$\Theta : \mathcal{D} \hookrightarrow L^2(\mu).$$

The adjoint makes “unstructured” $L^2(\mu)$ functions into holomorphic functions, so we expect it to be more manageable. Using the reproducing kernel property, we see that, for $g \in L^2(\mu)$

$$(19) \quad \begin{aligned} \Theta g(\zeta) &= \langle \Theta g, K_\zeta \rangle_{\mathcal{D}} \\ &= \langle g, K_\zeta \rangle_{L^2(\mu)} \\ &= \int_{\overline{\mathbb{D}}} g(z) K_z(\zeta) d\mu(z), \end{aligned}$$

because $\overline{K_\zeta(z)} = K_z(\zeta)$. We now insert (19) in the boundedness property of Θg :

$$\begin{aligned} C(\mu) \int_{\overline{\mathbb{D}}} |g|^2 d\mu &\geq \|\Theta g\|_{\mathcal{D}}^2 \\ &= \left\langle \int_{\overline{\mathbb{D}}} g(z) K_z(\cdot) d\mu(z), \int_{\overline{\mathbb{D}}} g(w) K_w(\cdot) d\mu(w) \right\rangle_{\mathcal{D}} \\ &= \int_{\overline{\mathbb{D}}} g(z) \int_{\overline{\mathbb{D}}} \overline{g(w)} d\mu(w) \langle K_z, K_w \rangle_{\mathcal{D}} d\mu(z) \\ &= \int_{\overline{\mathbb{D}}} g(z) \int_{\overline{\mathbb{D}}} \overline{g(w)} d\mu(w) K_z(w) d\mu(z). \end{aligned}$$

Overall, we have that the measure μ is Carleson for \mathcal{D} if and only if the weighted quadratic inequality

$$(20) \quad \int_{\mathbb{D}} g(z) \int_{\mathbb{D}} \overline{g(w)} d\mu(w) K_z(w) d\mu(z) \leq C(\mu) \int_{\mathbb{D}} |g|^2 d\mu$$

holds. Recalling that $k(z, w) = \Re K_z(w)$, it is clear that (20) implies

$$(21) \quad \int_{\mathbb{D}} g(z) \int_{\mathbb{D}} g(w) d\mu(w) k(z, w) d\mu(z) \leq C(\mu) \int_{\mathbb{D}} |g|^2 d\mu,$$

for real valued g and that, vice-versa, (21) for real valued g implies (20), with a twice larger constant: $\|\Theta(g_1 + ig_2)\|_{\mathcal{D}}^2 \leq 2(\|\Theta g_1\|_{\mathcal{D}}^2 + \|\Theta g_2\|_{\mathcal{D}}^2)$. [The same reasoning says that μ is Carleson if and only if (21) holds for positive g 's: the problem is reduced to a weighted inequality for a real (positive, in fact), symmetric kernel k]. Condition (17) is obtained by testing (21) over functions of the form $g = \chi_{\overline{S(\zeta)}}$. The finiteness of μ follows by testing the imbedding $\mathcal{D} \hookrightarrow L^2(\mu)$ on the function $f \equiv 1$.

The hard part is proving the sufficiency of (17): see [15], [10], [13], [34], [52] for different approaches to the problem. See also the very recent [58] for an approach covering the full range of the weighted Dirichlet spaces in the unit ball of \mathbb{C}^n . \square

The reasoning above works the same way with all reproducing kernels (provided the integrals involved make sense, of course). In particular, the problem of finding the Carleson measures for RKHS reduces, in general, to a weighted quadratic inequality like (21), with positive g 's.

3.2.1. *A Family of Necessary and Sufficient Testing Conditions.* Condition (4) is the endpoint of a family of such conditions, and the quadratic inequality (21) is the endpoint of a corresponding family of quadratic inequalities equivalent to the membership of μ to the Carleson class.

The kernels K and $k = \Re K$ define positive operators on \mathcal{D} , hence, By general Hilbert space theory, the boundedness inequality

$$\|\Theta g\|_{\mathcal{D}}^2 \leq C(\mu) \|g\|_{L^2(\mu)}^2$$

is equivalent to the boundedness of the operator $S : g \mapsto Sf = \int_{\mathbb{D}} k(\cdot, w) g(w) d\mu(w)$ on $L^2(\mu)$, i.e. to

$$(22) \quad \int_{\mathbb{D}} \left(\int_{\mathbb{D}} k(z, w) g(w) d\mu(w) \right)^2 d\mu(z) \leq C(\mu) \int_{\mathbb{D}} g^2 d\mu,$$

with the same constant $C(\mu)$. Testing (22) on $g = \chi_{\overline{S(\zeta)}}$ and restricting, we have the new testing condition

$$(23) \quad \int_{\overline{S(\zeta)}} \left(\int_{\overline{S(\zeta)}} k(z, w) d\mu(w) \right)^2 d\mu(z) \leq C(\mu) \mu(\overline{S(\zeta)}).$$

Observe that, by Jensen's inequality, (23) is *a priori* stronger than (4), although, by the preceding considerations, it is equivalent to it. Assuming the viewpoint that (23) represents the $L^2(\mu) - L^2(\mu)$ inequality for the "singular integral operator" having kernel ' k ', and using sophisticated machinery used to solve the Painlevé problem, Tchoundja [52] pushed this kind of analysis much further. Using also results in [10], he was able to prove the following.

Theorem 5 (Tchoundja, [52]). *Each of the following conditions on a finite measure μ is equivalent to the fact that $\mu \in CM(\mathcal{D})$:*

- For some $p \in (1, \infty)$ the following inequality holds,

$$(24) \quad \int_{\overline{\mathbb{D}}} \left(\int_{\overline{\mathbb{D}}} k(z, w) g(w) d\mu(w) \right)^p d\mu(z) \leq C_p(\mu) \int_{\overline{\mathbb{D}}} g^p d\mu.$$

- Inequality (24) holds for all $p \in (1, \infty)$.
- For some $p \in [1, \infty)$ the following testing condition holds,

$$(25) \quad \int_{\overline{S(\zeta)}} \left(\int_{\overline{S(\zeta)}} k(z, w) d\mu(w) \right)^p d\mu(z) \leq C_p(\mu) \mu(\overline{S(\zeta)}).$$

- The testing condition (25) holds for all $p \in [1, \infty)$.

Actually, Tchoundja deals with different spaces of holomorphic functions, but his results extend to the Dirichlet case. As mentioned earlier, the $p = 1$ endpoint of Theorem 5 is in [10].

3.2.2. *Another Family of Testing Conditions.* It was proved in [15] that a measure μ on $\overline{\mathbb{D}}$ is Carleson for \mathcal{D} if and only if (18) holds. In [36], Kerman and Sawyer had found another, seemingly weaker necessary and sufficient condition. In order to compare the two conditions, we restate (18) differently. Let $I(z) = \partial S(z) \cap \partial \mathbb{D}$. For $\theta \in I(z)$ and $s \in [0, 1 - |z|]$, let $S(\theta, s) = S((1 - s)e^{i\theta})$. Condition (18) is easily seen to be equivalent to have, for all $z \in \mathbb{D}$,

$$(26) \quad \int_{I(z)} \int_0^{1-|z|} \left(\frac{\mu(S(z) \cap \mu(S(\theta, s)))}{s^{1/2}} \right)^2 \frac{ds}{s} d\theta \leq C(\mu) \mu(S(z)).$$

Kerman and Sawyer proved that μ is a Carleson measure for \mathcal{D} if and only if, for all $z \in \mathbb{D}$,

$$(27) \quad \int_{I(z)} \sup_{s \in (0, 1-|z|]} \left(\frac{\mu(S(z) \cap \mu(S(\theta, s)))}{s^{1/2}} \right)^2 d\theta \leq C(\mu) \mu(S(z)).$$

Now, the quantity inside the integral on the left hand side of (27) is smaller than the corresponding quantity in (26). Due to the presence of the measure ds/s and the fact that the quantity $\mu(S(\theta, s))$ changes regularly with θ fixed and s variable; the domination of one by the other comes from the imbedding $\ell^2 \subseteq \ell^\infty$. The fact that, “on average”, the inclusion can be reversed is at first surprising. In fact, it is a consequence of the Muckenhoupt-Wheeden inequality [39] (or an extension of it), that the quantities on the left hand side of (26) and (27) are equivalent.

Theorem 6 ([36][15]). *A measure μ on $\overline{\mathbb{D}}$ is Carleson for the Dirichlet space \mathcal{D} if and only if it is finite and for some $p \in [1, \infty]$ (or, which is the same, for all $p \in [1, \infty]$) and all $z \in \mathbb{D}$:*

$$(28) \quad \int_{I(z)} \left[\int_0^{1-|z|} \left(\frac{\mu(S(z) \cap \mu(S(\theta, s)))}{s^{1/2}} \right)^p \right]^{2/p} \frac{ds}{s} d\theta \leq C(\mu) \mu(S(z)).$$

The inequality of Muckenhoupt and Wheeden was independently rediscovered by T. Wolff [32], with a completely new proof. Years later, trying to understand why the conditions in [15] and [36] were equivalent, although seemingly different, in [9] the authors, unaware of the results in [39] and [32], found another (direct) proof of the inequality.

4. THE TREE MODEL

4.1. The Bergman Tree. The unit disc \mathbb{D} can be discretized into Whitney boxes. The set of such boxes has a natural tree structure. In this section, we want to explain how analysis on the holomorphic Dirichlet space is related to analysis on similar spaces on the tree, and not only on a metaphoric level.

For integer $n \geq 0$ and $1 \leq k \leq 2^n$, consider the regions

$$\alpha(n, k) = \left\{ z = re^{i\theta} \in \mathbb{D} : 2^{-n} \leq 1 - |z| < 2^{-n-1}, \frac{k}{2^n} \leq \frac{\theta}{2\pi} < \frac{k+1}{2^n} \right\}.$$

We identify the set of regions with an abstract set indexing them. Let \mathcal{T} be the set of such regions. The elements of \mathcal{T} are a partition of the unit disc \mathbb{D} in regions whose Euclidean diameter, Euclidean in-radius and Euclidean distance to the boundary comparable to each other, with constants independent of the considered region. An easy exercise in hyperbolic geometry shows that the regions $\alpha \in \mathcal{T}$ have approximatively the same hyperbolic diameter and hyperbolic in-radius. We give the set \mathcal{T} two geometric-combinatorial structures: a *tree structure*, in which there is an edge between α and β when the corresponding regions share an arc of a circle; a *graph structure*, in which there is an edge between α and β if the closures of the corresponding regions have some point of \mathbb{D} in common. When referring to the graph structure, we write \mathcal{G} instead of \mathcal{T} .

In the tree \mathcal{T} we chose a distinguished point $o = \alpha(0, 1)$, the *root* of \mathcal{T} . The distance $d_{\mathcal{T}}(\alpha, \beta)$ between two points α, β in \mathcal{T} is the minimum number of edges of \mathcal{T} one has to travel going from the vertex α to the vertex β . Clearly, there is a unique path from α to β having minimal length: it is the *geodesic* $[\alpha, \beta]$ between α and β , which we consider as a set of points. The choice of the root gives \mathcal{T} a partial order structure: $\alpha \leq \beta$ if $\alpha \in [o, \beta]$. The *parent* of $\alpha \in \mathcal{T} \setminus \{o\}$ is the point α^{-1} on $[o, \alpha]$ such that $d(\alpha, \alpha^{-1}) = 1$. Each point α is the parent of two points in \mathcal{T} (its *children*), labeled when necessary as α^{\pm} . The natural geometry on \mathcal{T} is a simplified version of the hyperbolic geometry of the disc.

We might define a distance $d_{\mathcal{G}}$ on the graph \mathcal{G} using edges of \mathcal{G} instead of edges of \mathcal{T} . The distance $d_{\mathcal{G}}$ is realized by geodesics, although we do not have uniqueness anymore. (We have “almost uniqueness”: two geodesic between α and β maintain a reciprocal distance which is bounded by a positive constant C , independent of α and β). The following facts are rather easy to prove.

- (1) $d_{\mathcal{G}}(\alpha, \beta) \leq d_{\mathcal{T}}(\alpha, \beta)$.
- (2) If $z \in \alpha$ and $w \in \beta$, then $d_{\mathcal{G}}(\alpha, \beta) + 1 \approx d(z, w) + 1$: the graph metric is roughly the hyperbolic metric at unit scale.
- (3) There are points α_n, β_n such that $d_{\mathcal{T}}(\alpha_n, \beta_n)/d_{\mathcal{G}}(\alpha_n, \beta_n) \rightarrow \infty$: there are points which are close in the graph, but far away in the tree.

While the graph geometry is a good approximation of the hyperbolic geometry at a fixed scale, the same can not be said about the tree geometry. Nonetheless, the tree geometry is much more elementary, and it is that we are going to use. It is a bit surprising that, in spite of the distortion of the hyperbolic metric pointed out in (3), the tree geometry is so useful.

Let us introduce the analogs of cones and Carleson boxes on the tree: $\mathcal{P}(\alpha) = [o, \alpha] \subset \mathcal{T}$ is the *predecessor set* of $\alpha \in \mathcal{T}$ (when you try to picture it, you get a sort of cone) and $\mathcal{S}(\alpha) = \{\beta \in \mathcal{T} : \alpha \in \mathcal{P}(\beta)\}$, its dual object, is the *successor set* of α (a sort of Carleson box).

Given $\alpha, \beta \in \mathcal{T}$, we denote their *confluent* by $\alpha \wedge \beta$: the point on the geodesic between α and β which is closest to the root o . That is,

$$\mathcal{P}(\alpha \wedge \beta) = \mathcal{P}(\alpha) \cap \mathcal{P}(\beta).$$

In terms of \mathbb{D} geometry, the confluent corresponds to the highest point of the smallest Carleson box containing two points; if $z, w \in \mathbb{D}$ are the points, the point which plays the rôle of $\alpha \wedge \beta$ is roughly the point having argument halfway between that of z and that of w , and having Euclidean distance $|1 - z\bar{w}|$ from the boundary.

4.2. Detour: The Boundary of the Tree and its Relation with the Disc's Boundary. The distortion of the metric induced by the tree structure has an interesting effect on the boundary. One can define a boundary $\partial\mathcal{T}$ of the tree \mathcal{T} . While the boundary of \mathbb{D} (which we might think of as a boundary for the graph \mathcal{G}) is connected, the boundary $\partial\mathcal{T}$ is totally disconnected; it is in fact homeomorphic to a Cantor set. Notions of boundaries for graphs, and trees in particular, are an old a useful topic in probability and potential theory. We mention [46] as a nice introduction to this topic.

We will see promptly that the boundary $\partial\mathcal{T}$ is compact with respect to a natural metric and that, as such, it carries positive Borel measures. Furthermore, if μ is positive, Borel measure without atoms with support on $\partial\mathbb{D}$, then it can be identified with a positive, Borel measure without atoms on $\partial\mathcal{T}$.

This is the main reason we are interested in trees and tree's boundary. Some theorems are easier to prove on the tree's boundary, some estimates become more transparent and some objects are easier to picture. Often, it is possible to split a problem in two parts: a “soft” part, to deal with in the disc geometry, and a “hard” combinatorial part, which one can formulate and solve in the easier tree geometry. Many of these results and objects can then be transplanted in the context of the Dirichlet space.

As a set, the boundary $\partial\mathcal{T}$ contains as elements the half-infinite geodesics on \mathcal{T} , having o as endpoint. For convenience, we think of $\zeta \in \partial\mathcal{T}$ as of a point and we denote by $\mathcal{P}(\zeta) = [o, \zeta) \subset \mathcal{T}$ the geodesic labeled by ζ . We introduce on $\partial\mathcal{T}$ a metric which mimics the Euclidean metric on the disc:

$$\delta_{\mathcal{T}}(\zeta, \xi) = 2^{-d_{\mathcal{T}}(\zeta \wedge \xi)},$$

where $\zeta \wedge \xi$ is defined as in the “finite” case $\alpha, \beta \in \mathcal{T}$: $\mathcal{P}(\zeta \wedge \xi) = \mathcal{P}(\zeta) \cap \mathcal{P}(\xi)$. It is easily verified that, modulo a multiplicative constant, $\delta_{\mathcal{T}}$ is the weighted length of the doubly infinite geodesic $\gamma(\zeta, \xi)$ which joins ζ and ξ , where the weight assigns to each edge $[\alpha, \alpha^{-1}]$ the number $2^{-d_{\mathcal{T}}(\alpha)}$. The metric can be extended to $\overline{\mathcal{T}} = \mathcal{T} \cup \partial\mathcal{T}$ by similarly measuring geodesics' lengths for all geodesics. This way, we obtain a compact metric space $(\overline{\mathcal{T}}, \delta_{\mathcal{T}})$, where \mathcal{T} is a discrete subset of $\overline{\mathcal{T}}$, having $\partial\mathcal{T}$ as metric boundary. *The subset $\partial\mathcal{T}$, as we said before, turns out to be a totally disconnected, perfect set.*

The relationship between $\partial\mathcal{T}$ and $\partial\mathbb{D}$ is more than metaphoric. For a point $\zeta \in \partial\mathcal{T}$, let $\mathcal{P}(\zeta) = \{\zeta_n : n \in \mathbb{N}\}$ be an enumeration of the points $\zeta_n \in \mathcal{T}$ of the corresponding geodesic, ordered in such a way that $d(\zeta_n, o) = n$. Each ζ in \mathcal{T} can be identified with a dyadic sub-arc of $\partial\mathbb{D}$: if $Q(\zeta)$ is the Whitney box labeled by ζ , $\zeta \leftrightarrow I(\zeta) = \partial Q(\zeta) \cap \partial\mathbb{D}$. Define the map $\Lambda : \partial\mathcal{T} \rightarrow \partial\mathbb{D}$,

$$(29) \quad \Lambda(\zeta) = \bigcap_{n \in \mathbb{N}} I(\zeta_n).$$

It is easily verified that Λ is a Lipschitz continuous map of $\partial\mathcal{T}$ onto $\partial\mathbb{D}$, which fails to be injective at a countable set (the set of the dyadic rationals $\times 2\pi$). More important is the (elementary, but not obvious) fact that Λ maps Borel measurable sets in $\partial\mathcal{T}$ to Borel measurable sets in $\partial\mathbb{D}$. This allows us to move Borel measures back and forth from $\partial\mathcal{T}$ to $\partial\mathbb{D}$.

Given a Borel, positive measure ω on $\partial\mathcal{T}$, let $(\Lambda_*\omega)(E) = \omega(\Lambda^{-1}(E))$ be the usual push-forward measure. Given a positive, Borel measure μ on \mathbb{S} , define its **pull-back** $\Lambda^*\mu$ to be the positive Borel measure on $\partial\mathcal{T}$

$$(30) \quad (\Lambda^*\mu)(F) = \int_{\mathbb{S}} \frac{\sharp(\Lambda^{-1}(\theta) \cap A)}{\sharp(\Lambda^{-1}(\theta))} d\mu(e^{i\theta}).$$

Proposition 7. (i) *The integrand in (30) is measurable, hence the integral is well-defined.*

(ii) $\Lambda_*(\Lambda^*(\mu)) = \mu$.

(iii) *For any closed subset A of \mathbb{S} , $\Lambda_*\omega(A) = \omega(\Lambda^{-1}(A))$, by definition.*

(iv) *For any closed subset B of $\partial\mathcal{T}$, $\Lambda^*\mu(B) \approx \mu(\Lambda(B))$.*

(v) *In (iv), we have equality if the measure μ has no atoms.*

See [18] for more general versions of the proposition.

4.3. A Version of the Dirichlet Space on the Tree. Consider the Hardy-type operator \mathcal{I} acting on functions $\varphi : \mathcal{T} \rightarrow \mathbb{R}$,

$$\mathcal{I}\varphi(\alpha) = \sum_{\beta \in \mathcal{P}(\alpha)} \varphi(\beta).$$

The Dirichlet space $\mathcal{D}_{\mathcal{T}}$ on \mathcal{T} is the space of the functions $\Phi = \mathcal{I}\varphi$, $\varphi \in \ell^2(\mathcal{T})$, with norm $\|\Phi\|_{\mathcal{D}_{\mathcal{T}}} = \|\varphi\|_{\ell^2}$. Actually, we will always talk about the space ℓ^2 and the operator \mathcal{I} , rather than about the space $\mathcal{D}_{\mathcal{T}}$, which is however the *trait d'union* between the discrete and the continuous theory.

What we are thinking of, in fact, is discretizing a Dirichlet function $f \in \mathcal{D}$ in such a way

- (1) $\varphi(\alpha) \sim (1 - |z(\alpha)|)|f'(z(\alpha))|$, where $z(\alpha)$ is a distinguished point in the region α (or in its closure);
- (2) $\mathcal{I}\varphi(\alpha) = f(\alpha)$.

Let us mention a simple example from [14], saying that ℓ^2 is “larger” than \mathcal{D} .

Proposition 8. *Consider a subset $\{z(\alpha) : \alpha \in \mathcal{T}\}$ of \mathbb{D} , where $z(\alpha) \in \alpha$, and let $f \in \mathcal{D}$. Then, there is a function φ in $\ell^2(\mathcal{T})$ such that $\mathcal{I}\varphi(\alpha) = f(z(\alpha))$ for all $\alpha \in \mathcal{T}$ and $\|\varphi\|_{\ell^2} \lesssim \|f\|_{\mathcal{D}}$.*

Proof. Assume without loss of generality that $f(0) = 0$ and let $\varphi(\alpha) := f(z(\alpha)) - f(z(\alpha^{-1}))$. By telescoping, $\varphi(\alpha) = f(z(\alpha))$. To prove the estimate,

$$\begin{aligned} \|h\|_{\ell^2(\mathcal{T})}^p &= \sum_{\alpha} |f(z(\alpha)) - f(z(\alpha^{-1}))|^2 \\ &\lesssim \sum_{\alpha} |(1 - |z(\alpha)|)f'(w(\alpha))|^2 \\ &\quad \text{for some } w(\alpha) \text{ in the closure of } \alpha, \\ &\approx \sum_{\alpha} (1 - |z(\alpha)|)^2 \left| \frac{1}{(1 - |z_{\alpha}|)^2} \int_{\zeta \in \mathbb{D}: |\zeta - w(\alpha)| \leq (1 - |z(\alpha)|)/10} f'(\zeta) dA(\zeta) \right|^2 \\ &\quad \text{by the (local) Mean Value Property,} \end{aligned}$$

$$\begin{aligned}
&\lesssim \sum_{\alpha} \int_{\zeta: |\zeta - w(\alpha)| \leq (1 - |z(\alpha)|)/10} |f'(\zeta)|^2 dA(\zeta) \\
&\quad \text{by Jensen's inequality,} \\
&\approx \|f\|_{\mathcal{D}}^2,
\end{aligned}$$

since the discs $\{\zeta : |\zeta - w(\alpha)| \leq (1 - |z(\alpha)|)/10\}$ clearly have bounded overlapping. \square

4.4. Carleson Measures and All That, on the Tree and on the Disc. Let μ be a positive measure on the unit disc. Identify it with a positive measure on \mathcal{T} , letting

$$\mu(\alpha) = \int_{\alpha} d\mu(\alpha).$$

4.4.1. Carleson Measures. We say that μ is a *Carleson measure* for $\mathcal{D}_{\mathcal{T}}$ if the operator $\mathcal{I} : \ell^2(\mathcal{T}) \rightarrow \ell^2(\mathcal{T}, \mu)$ is bounded. We write $\mu \in CM(\mathcal{D}_{\mathcal{T}})$.

Theorem 9. *We have that $CM(\mathcal{D}) = CM(\mathcal{D}_{\mathcal{T}})$, with comparable norms.*

Proof discussion. We can use the restriction argument of Proposition 8 to show that $CM(\mathcal{T}) \subseteq CM(\mathcal{D})$. Suppose for simplicity that $\mu(\partial\mathbb{D}) = 0$ (dealing with this case requires further discussion of the tree's boundary) and that $\mu \in CM(\mathcal{T})$:

$$\begin{aligned}
\int_{\mathbb{D}} |f|^2 d\mu &= \sum_{\alpha} \int_{\alpha} |f|^2 d\mu \leq \sum_{\alpha} \mu(\alpha) |f(z(\alpha))|^2 \\
&\quad \text{for some } z(\alpha) \text{ on the boundary of } \alpha \\
&= \sum_{\alpha} \mathcal{I}\varphi(\alpha) \mu(\alpha) \\
&\quad \text{with } \varphi \text{ as in Proposition 8} \\
&\leq \|\varphi\|_{\ell^2(\mathcal{T})}^2,
\end{aligned}$$

which proves the inclusion.

In the other direction, we use the duality argument used in the proof of Theorem 4. The fact that μ is Carleson for \mathcal{D} is equivalent to the boundedness of Θ , the adjoint of the imbedding, and this is equivalent to the inequality

$$\begin{aligned}
C(\mu) \int_{\mathbb{D}} |g|^2 d\mu &\geq \|\Theta g\|_{\mathcal{D}}^2 = \int_{\mathbb{D}} |(\Theta g)'(z)|^2 dA(z) \\
&\quad \text{this time we use a different way to compute the norm,} \\
&\geq \int_{\mathbb{D}} \left| \int_{\mathbb{D}} \frac{d}{dz} K(z, w) g(w) d\mu(w) \right|^2 dA(z) \\
(31) \quad &= \int_{\mathbb{D}} \left| \int_{\mathbb{D}} \frac{\bar{w}}{1 - \bar{w}z} g(w) d\mu(w) \right|^2 dA(z).
\end{aligned}$$

Testing (31) over all functions $g(w) = h(w)/\bar{w}$ with $h \geq 0$ and using the geometric properties of the kernel's derivative, we see that

$$(32) \quad C(\mu) \int_{\mathbb{D}} |g|^2 d\mu \geq \int_{\mathbb{D}} \left| \int_{S(z)} \bar{w} g(w) d\mu(w) \right|^2 dA(z).$$

We can further restrict to the case where h is constant on Whitney boxes ($h = \sum_{\alpha \in \mathcal{T}} \psi(\alpha) \chi_{\alpha}$) and, further restricting the integral, we see that (32) reduces to

$$(33) \quad C(\mu) \|\psi\|_{\ell^2(\mu)}^2 \geq \|\mathcal{I}^*(\psi d\mu)\|_{\ell^2}^2.$$

A duality argument similar (in the converse direction) to the previous one, this time in tree-based function spaces, shows that the last assertion is equivalent to having $\mathcal{I} : \ell^2(\mathcal{T}) \rightarrow \ell^2(\mathcal{T}, \mu)$ bounded, i.e. $\mu \in CM(\mathcal{T})$. \square

The proof could be carried out in the dual side, completely. Actually, this is almost obliged in several extensions of the theorem (to higher dimensions [14], to “sub-diagonal” couple of indices [8], etcetera). A critical analysis of the proof and some further considerations about the boundary of the tree show that Carleson measures satisfy a stronger property.

Corollary 10 (Arcozzi-Rochberg-Sawyer, [13]). *Let*

$$V(f)(Re^{i\theta}) = \int_0^R |f'(re^{i\theta})| dr$$

be the radial variation of $f \in \mathcal{D}$ (i.e. the length of the image of the radius $[0, Re^{i\theta}]$ under f). Then, $\mu \in CM(\mathcal{D})$ if and only if the stronger inequality

$$\int_{\mathbb{D}} V(f)^2 d\mu \leq C(\mu) \|f\|_{\mathcal{D}}^2.$$

holds.

Indeed, this remark is meaningful when μ is supported on $\partial\mathbb{D}$.

4.4.2. *Testing Conditions in the Tree Language.* In the proof discussion following Theorem 9, we ended by showing that a necessary and sufficient condition for a measure μ on \mathbb{D} to be in $CM(\mathbb{D})$ is (33). Making duality explicit, one computes

$$\mathcal{I}^*(\psi d\mu)(\alpha) = \int_{\overline{\mathcal{S}(\alpha)}} g d\mu.$$

Using as testing functions $g = \chi_{\overline{\mathcal{S}(\alpha_0)}}$, $\alpha_0 \in \mathcal{T}$ and throwing away some terms on the right hand side, we obtain the discrete testing condition:

$$(34) \quad C(\mu)\mu(\overline{\mathcal{S}(\alpha_0)}) \geq \sum_{\alpha \in \mathcal{S}(\alpha_0)} [\mu(\overline{\mathcal{S}(\alpha)})]^2.$$

We will denote by $[\mu]$ the best constant in (34).

Theorem 11 (Arcozzi-Rochberg-Sawyer, [15]). *A measure μ on \mathbb{D} belongs to $CM(\mathcal{D})$ if, and only if, it is finite and it satisfies (34).*

Given Theorem 9, Theorem 11 really becomes a characterization of the weighted inequalities for the operator \mathcal{I} (and/or its adjoint). There is a vast literature on weighted inequalities for operators having positive kernels, and virtually all of the proofs translate in the present context. Theorem 11 was proved in [15] by means of a *good- λ* argument. A different proof could be deduced by the methods in [34], where a deep equivalence is established between weighted inequalities and a class of integral (nonlinear) equations. In [13] a very short proof is given in terms of a maximal inequality.

The fact that the (discrete) testing condition (34) characterizes Carleson measures raises two natural questions.

- Is there a *direct* proof that the testing condition (34) is equivalent to Stegenga’s capacity condition?

- Is there an “explanation” of how a condition which is expressed in terms of the *tree structure* is sufficient to characterize properties whose natural environment is the *graph structure* of the unit disc?

4.4.3. *Capacities on the Tree.* Let E be a closed subset of $\partial\mathcal{T}$. We define a logarithmic-type and a Bessel-type capacity for E . As in the continuous case, they turn out to be equivalent.

The operator \mathcal{I} can be extended in the obvious way on the boundary of the tree, $\mathcal{I}\varphi(\zeta) = \sum_{\beta \in \mathcal{P}(\zeta)} \varphi(\beta)$ for ζ in $\partial\mathcal{T}$.

$$(35) \quad \text{Cap}_{\mathcal{T}}(E) = \inf \left\{ \|\varphi\|_{\ell^2(\mathcal{T})}^2 : \mathcal{I}\varphi(\zeta) \geq 1 \text{ on } E \right\}$$

will be the *tree capacity* of E , which roughly corresponds to logarithmic capacity.

Define the kernel $k_{\partial\mathcal{T}} : \partial\mathcal{T} \times \partial\mathcal{T} \rightarrow [0, +\infty]$,

$$k_{\partial\mathcal{T}}(\zeta, \xi) = 2^{d_{\mathcal{T}}(\zeta \wedge \xi)/2},$$

which mimics the Bessel kernel $k_{\mathbb{S}, 1/2}$. The energy of a measure ω on $\partial\mathcal{T}$ associated with the kernel is

$$\mathcal{E}_{\partial\mathcal{T}}(\omega) = \int_{\partial\mathcal{T}} (k_{\partial\mathcal{T}}\omega(\zeta))^2 dm_{\partial\mathcal{T}}(\zeta),$$

where $m_{\partial\mathcal{T}} = \Lambda^*m$ is the pullback of the linear measure on \mathbb{S} . More concretely, $m_{\partial\mathcal{T}}\partial\mathcal{S}(\alpha) = 2^{-d_{\mathcal{T}}(\alpha)}$. We define another capacity

$$\text{Cap}_{\partial\mathcal{T}}(E) = \sup \left\{ \frac{\omega(E)}{\mathcal{E}(\omega)} : \text{supp}(\omega) \subseteq E \right\},$$

the supremum being taken over positive, Borel measures on $\partial\mathcal{T}$.

As in the continuous case (with a simpler proof) one has that the two capacities are equivalent,

$$\text{Cap}_{\mathcal{T}}(E) \approx \text{Cap}_{\partial\mathcal{T}}(E).$$

It is not obvious that both are equivalent to the logarithmic capacity.

Theorem 12 (Benjamini-Peres, [22]).

$$\text{Cap}_{\mathcal{T}}(E) \approx \text{Cap}_{\partial\mathcal{T}}(E) \approx \text{Cap}_{\mathbb{S}, 1/2}(\Lambda(E)).$$

See [18] for an extension of this result to Bessel-type capacities on Ahlfors-regular metric spaces.

Proof discussion. Let ω be a positive Borel measure on $\partial\mathcal{T}$ and μ be a positive Borel measure on \mathbb{S} . It suffices to show that the energy of ω with respect to the kernel $k_{\partial\mathcal{T}}$ is comparable with the energy of $\Lambda_*\omega$ with respect to $k_{\mathbb{S}, 1/2}$ and that the energy of μ is comparable with energy of $\Lambda^*\mu$, with respect to the same kernels, obviously taken in reverse order. We can also assume the measures to be atomless, since atoms, both in \mathbb{S} and $\partial\mathcal{T}$, have infinite energy. Proposition 29 implies that the measure $\Lambda^*\mu$ is well defined and helps with the energy estimates, which are rather elementary. \square

Theorem 12 has direct applications to the theory of the Dirichlet space.

- As explained in [18], there is a direct relationship between tree capacity $\text{Cap}_{\mathcal{T}}$ and Carleson measures for the Dirichlet space. Namely, for a closed subset E of $\partial\mathcal{T}$,

$$(36) \quad \text{Cap}_{\mathcal{T}}(E) = \sup_{\mu: \text{supp}(\mu) \subseteq E} \frac{\mu(E)}{[\mu]}.$$

- As a consequence, we have that sets having null capacity are exactly sets which do not support positive Carleson measures. Together with Corollary (10) and the theorem of Benjamini and Peres, this fact implies an old theorem by Beurling.

Theorem 13 (Beurling, [20]).

$$\text{Cap}_{\mathbb{S},1/2}(\{\zeta \in \mathbb{S} : V(f)(e^{i\theta}) = +\infty\}) = 0.$$

In particular, Dirichlet functions have boundary values at all points on \mathbb{S} , but for a subset having null capacity. This result, the basis for the study of boundary behavior of Dirichlet functions, explains the differences and similarities between Hardy and Dirichlet theories. It makes it clear that capacity is for \mathcal{D} what arclength measure is in H^2 . On the other hand, there are Hardy functions (even bounded analytic functions) having infinite radial variation at almost all points on \mathbb{S} : radial variation is for the most part a peculiarly Dirichlet topic.

- Another application is in [16], where boundedness of certain bilinear forms on \mathcal{D} is discussed (and which also contains a different proof of Theorem [22], of which we were not aware at the moment of writing the article). Central to the proof of the main result is the holomorphic approximation of the discrete potentials which are extremal for the tree capacity of certain sets. See Section 6 for a discussion of this and related topics.

4.4.4. *Capacitary Conditions and Testing Conditions.* The capacitary condition of Stegenga and the discrete testing condition (34) (plus boundedness of μ) are equivalent, since both characterize $CM(\mathcal{D})$. It is easy to see that the capacitary condition is *a priori* stronger than the testing condition. A *direct* proof that the testing condition implies the capacitary condition is in [11]. The main tool in the proof is the characterization (36) of the tree capacity.

5. THE COMPLETE NEVANLINNA-PICK PROPERTY

In 1916 Georg Pick published the solution to the following interpolation problem.

Problem 14. *Given domain points $\{z_i\}_{i=1}^n \subset \mathbb{D}$ and target points $\{w_i\}_{i=1}^n \subset \mathbb{D}$ what is a necessary and sufficient condition for there to be an $f \in H^\infty$, $\|f\|_\infty \leq 1$ which solves the interpolation problem $f(z_i) = w_i$ $i = 1, \dots, n$?*

A few years later Rolf Nevanlinna independently found an alternative solution. The problem is now sometimes called Pick's problem and sometimes goes with both names; Pick-Nevanlinna (chronological) and Nevanlinna-Pick (alphabetical). The result has been extraordinarily influential and many of the topics in this volume have intellectual roots in that problem.

One modern extension of Pick's question is the following:

Problem 15 (Pick Interpolation Question). *Suppose H is a Hilbert space of holomorphic functions on \mathbb{D} . Given $\{z_i\}_{i=1}^n, \{w_i\}_{i=1}^n \subset \mathbb{D}$ is there a function m in \mathcal{M}_H , the multiplier algebra, with $\|m\|_{\mathcal{M}_H} \leq 1$, which performs the interpolation $m(z_i) = w_i; i = 1, 2, \dots, n$*

There is a necessary condition for the interpolation problem to have a solution which holds for any RKHS. We develop that now.

Suppose we are given the data for the interpolation question. Let V be the span of the kernel functions $\{k_i\}_{i=1}^n$.

Theorem 16. *Define the map T of that V to itself by*

$$T\left(\sum a_i k_i\right) = \sum a_i \bar{w}_i k_i.$$

A necessary condition for the Pick Interpolation Question to have a positive answer is that $\|T\| \leq 1$. Equivalently a necessary condition is that the associated matrix

$$(37) \quad \text{Mx}(T) = \left((1 - w_j \bar{w}_i) k_j(z_i)\right)_{i,j=1}^n$$

be positive semi-definite; $\text{Mx}(T) \geq 0$.

Proof. Suppose there is such a multiplier m and let M be the operator of multiplication by m acting on H . We have $\|M\| = \|m\|_{\mathcal{M}(H)} \leq 1$. Hence the adjoint operator, M^* satisfies $\|M^*\| \leq 1$. We know that given $\zeta \in \mathbb{D}$, $M^*k_\zeta = \overline{m(\zeta)}k_\zeta$. Thus V is an invariant subspace for M^* and the restriction of M^* to V is the operator T of the theorem. Also the restriction of M^* to V has, *a fortiori*, norm at most one. That gives the first statement.

The fact that the norm of T is at most one means that for scalars $\{a_i\}_{i=1}^n$ we have

$$\left\|\sum a_i \bar{w}_i k_i\right\|^2 \leq \left\|\sum a_i k_i\right\|^2.$$

We compute the norms explicitly recalling that $\langle k_i, k_j \rangle = k_i(z_j)$ and rearrange the terms and find that

$$\sum_{i,j} (1 - w_j \bar{w}_i) k_j(z_i) a_j \bar{a}_i \geq 0.$$

The scalars $\{a_i\}_{i=1}^n$ were arbitrary and thus this is the condition that $\text{Mx}(T) \geq 0$. \square

The matrix $\text{Mx}(T)$ is called the *Pick matrix* of the problem. For the Hardy space it takes the form

$$\text{Mx}(T) = \left(\frac{1 - w_i \bar{w}_j}{1 - z_i \bar{z}_j}\right)_{i,j=1}^n$$

Theorem 17 (Pick). *For the Hardy space, the necessary condition for the interpolation problem to have a solution, (37), is also sufficient.*

See [2] for a proof.

Remark 18. *The analog of Pick's theorem fails for the Bergman space; (37) is not sufficient.*

It is now understood that there are classes of RKHS for which the condition (37) is sufficient for the interpolation problem to have a solution. Such spaces are said to have the Pick property. In fact there is a subclass, those with the Complete Pick Property, CNPP for which (37) is a sufficient condition for the interpolation

problem to have a solution and there is also a matricial analog of (37) is a necessary and sufficient for a matricial analog of the interpolation problem to have a solution.

It is a consequence of the general theory of spaces with the CNPP that the kernel functions never vanish; $\forall z, w \in X, k_z(w) \neq 0$.

For spaces of the type we are considering there is a surprisingly simple characterization of spaces with the CNPP. Suppose H is a Hilbert space of holomorphic functions on the disk in which the monomials $\{z^n\}_{n=0}^\infty$ are a complete orthogonal set. The argument we used to identify the reproducing kernel for the Dirichlet space can be used again and we find that for $\zeta \in \mathbb{D}$ we have

$$\begin{aligned} k_\zeta^H(z) &= \sum_{n=0}^{\infty} \frac{\bar{\zeta}^n z^n}{\|z^n\|_H^2} \\ &= \sum_{n=0}^{\infty} a_n \bar{\zeta}^n z^n \end{aligned}$$

We know that $a_0 = \|1\|_H^{-2} > 0$ hence in a neighborhood of the origin the function $\sum_{n=0}^{\infty} a_n t^n$ has a reciprocal given by a power series. Define $\{c_n\}$ by

$$(38) \quad \frac{1}{\sum_{n=0}^{\infty} a_n t^n} = \sum_{n=0}^{\infty} c_n t^n.$$

Having $a_0 > 0$ insures $c_0 > 0$.

Theorem 19. *The space H has the CNPP if and only if*

$$c_n \leq 0 \quad \forall n > 0.$$

Using this we immediately see that the Hardy space has the CNPP and the Bergman space does not.

Theorem 20. *The Dirichlet space \mathcal{D} with the norm $\|\cdot\|_{\mathcal{D}}$,*

$$\left\| \sum_{n=0}^{\infty} b_n z^n \right\|_{\mathcal{D}}^2 = \sum_{n=0}^{\infty} (n+1) |b_n|^2,$$

has the complete Nevanlinna-Pick property.

On the other hand one needs only compute a few of the c_n to find out that:

Remark 21. *The space \mathcal{D} with the norm*

$$\left\| \sum_{n=0}^{\infty} b_n z^n \right\|_{\mathcal{D}}^2 = |b_0|^2 + \sum_{n=1}^{\infty} n |b_n|^2$$

does not have the CNPP.

If a RKHS has the CNPP then a number of other subtle and interesting consequences follow. In particular, this applies for the Dirichlet space.

We refer the reader to the foundational article [3] and to the beautiful monograph [2] for a comprehensive introduction to spaces with the CNPP.

6. THE MULTIPLIER SPACE AND OTHER SPACES INTRINSIC TO \mathcal{D} THEORY

6.1. **Multipliers.** Suppose H is a RKHS of holomorphic functions in the disk. We say that a function m is a *multiplier* (of H or for H) if multiplication by m maps H boundedly to itself; that is there is a $C = C(m)$ so that for all $h \in H$

$$\|mh\|_H \leq C \|h\|_H.$$

Let \mathcal{M}_H be the space of all multipliers of H and for $m \in \mathcal{M}_H$ let $\|m\|_{\mathcal{M}_H}$ be the operator norm of the multiplication operator. With this norm \mathcal{M}_H is a commutative Banach algebra.

It is sometimes easy and sometimes difficult to get a complete description of the multipliers of a given space H . If the constant functions are in H (they are, in the case of the Hardy and of the Dirichlet space), then $\mathcal{M}_H \subset H$. In fact for $\|1\|_H = 1$ and hence the inclusion is contractive: $\|m\|_H = \|m \cdot 1\|_{\mathcal{M}_H} \leq \|m\|_{\mathcal{M}_H} \|1\|_H = \|m\|_{\mathcal{M}_H}$.

Also, for each of \mathcal{D} , H^2 , and A^2 (the Bergman space) the multiplier algebra is contractively contained in H^∞ ,

$$\|m\|_{H^\infty} \leq \|m\|_{\mathcal{M}_H}.$$

One way to see this is by looking at the action of the adjoint of the multiplication operator on reproducing kernels. Let H be one of \mathcal{D} , H^2 , and A^2 ; let $m \in \mathcal{M}_H$ and let M be the operator of multiplication by m acting on H . Let M^* be the adjoint of the operator M . We select $\zeta, z \in \mathbb{D}$ and compute

$$\begin{aligned} M^*k_{H,\zeta}(z) &= \langle M^*k_{H,\zeta}, k_{H,z} \rangle \\ &= \langle k_{H,\zeta}, Mk_{H,z} \rangle \\ &= \langle k_{H,\zeta}, mk_{H,z} \rangle \\ &= \overline{\langle mk_{H,z}, k_{H,\zeta} \rangle} \\ &= \overline{m(\zeta)k_{H,z}(\zeta)} \\ &= \overline{m(\zeta)}k_{H,\zeta}(z). \end{aligned}$$

Thus $k_{H,\zeta}$ is an eigenvector of M^* , the adjoint of the multiplication operator, with eigenvalue $\overline{m(\zeta)}$. Hence $|m(\zeta)| \leq \|M^*\| = \|M\|$. Taking the supremum over $\zeta \in \mathbb{D}$ gives the desired estimate.

For the Hardy space that is the full story; $\mathcal{M}_{H^2} = H^\infty$. In the Dirichlet case, things are a bit more complicated.

Proposition 22. *m is a multiplier for the Dirichlet space if and only if $m \in H^\infty$ and $d\mu_m(z) = |m'(z)|^2 dx dy \in CM(\mathcal{D})$.*

This was the motivation for Stegenga's study [50] of the Carleson measures for the Dirichlet space. Observe that $\int_{\mathbb{D}} d\mu_m = \|m\|_{\mathcal{D},*}^2$.

Let us look again at the Hardy case, in the light of Stegenga's Proposition 22. Let χ_{H^2} be the space of the functions m holomorphic in \mathbb{D} such that the measure $d\lambda_m(z) = (1 - |z|^2)|m'(z)|^2 dA(z)$ is a Carleson measure for the Hardy space. The reason for choosing such measure is that $\int_{\mathbb{D}} d\lambda_m(z) \approx \|m\|_{H^2}^2$ (if $m(0) = 0$), as in the Dirichlet case. Now, it is known that $\chi_{H^2} = BMOA$ is the space of the analytic functions in BMO . Proposition 22 says that multiplier algebra of \mathcal{D} is exactly $\chi \cap H^\infty$ (here, χ contains the functions m such that $d\mu_m(z) = |m'(z)|^2 dx dy \in CM(\mathcal{D})$). The analogous result for H^2 would be: the multiplier space of H^2 consists

of the functions in $BMOA$ which are essentially bounded. This is true, but not very interesting, since $H^\infty \subseteq BMOA$.

6.2. The Weakly Factored Space $\mathcal{D} \odot \mathcal{D}$ and its Dual.

6.2.1. *Some Facts from H^2 Theory.* It is well known that the some spaces of holomorphic functions naturally arise within H^2 theory: H^1 , H^∞ , BMO . We shortly recall some of their mutual connections. We have just seen that H^∞ naturally arises as the multiplier algebra of H^2 : $Mult(H^2) = H^\infty$. On the other hand, by the inner/outer factorization of H^2 functions it easily follows that $H^1 = H^2 \cdot H^2$ is the space of products of H^2 functions. C. Fefferman's celebrated theorem says that $(H^1)^* = BMO$ is the space of analytic functions with bounded mean oscillation. Functions in BMO are defined by the well-know, elegant integral property to which they owe their name, but could be otherwise defined as the functions b analytic in \mathbb{D} such that $d\mu_b = (1 - |z|^2)|b'(z)|^2 dA(z)$ is a Carleson measure for H^2 :

$$\int_{\mathbb{D}} |f|^2 d\mu_b \leq C(\mu) \|f\|_{H^2}^2.$$

It might be added that the multiplier algebra H^∞ consists of those functions in BMO which are bounded, but this is never done, since the multiplier algebra *coincides* with H^∞ .

The spaces just considered are linked with the Hankel forms and Nehari's Theorem. Given analytic b , define the *Hankel form* with symbol b as

$$T_b(f, g) = \langle b, fg \rangle_{H^2}.$$

It was shown by Nehari that

$$\sup_{f, g \in H^2} \frac{|T_b(f, g)|}{\|f\|_{H^2} \|g\|_{H^2}} \approx \|b\|_{(H^1)^*} \approx \|b\|_{BMO},$$

the last equality following from Fefferman's Theorem.

6.2.2. *Function Spaces Naturally Related with the Dirichlet Space.* One might first think that, being the Dirichlet space naturally defined in terms of hyperbolic geometry, the spaces playing the rôle of H^1 , H^∞ and BMO in Dirichlet theory would be the Bloch space \mathcal{B} , defined by the (conformally invariant) norm:

$$\|f\|_{\mathcal{B}} = \|\delta f\|_{L^\infty(\mathbb{D})} = \sup_{z \in \mathbb{D}} (1 - |z|^2) |f'(z)|$$

and similarly defined invariant spaces. It turns out that, from the viewpoint Hilbert space function theory, the relevant spaces are others.

Define the weakly factored space $\mathcal{D} \odot \mathcal{D}$ to be the completion of finite sums $h = \sum f_j g_j$ using the norm

$$\|h\|_{\mathcal{D} \odot \mathcal{D}} = \inf \left\{ \sum \|f_j\|_{\mathcal{D}} \|g_j\|_{\mathcal{D}} : h = \sum f_j g_j \right\}.$$

In particular if $f \in \mathcal{D}$ then $f^2 \in \mathcal{D} \odot \mathcal{D}$ and

$$(39) \quad \|f^2\|_{\mathcal{D} \odot \mathcal{D}} \leq \|f\|_{\mathcal{D}}^2.$$

It is immediate that, in the Hardy case, $H^2 \odot H^2 = H^2 \cdot H^2 = H^1$.

We also introduce a variant of $\mathcal{D} \odot \mathcal{D}$. Define the space $\partial^{-1}(\partial\mathcal{D} \odot \mathcal{D})$ to be the completion of the space of functions h such that h' can be written as a finite sum, $h' = \sum f'_j g_j$ (and thus $h = \partial^{-1} \sum (\partial f_i) g_i$), with the norm

$$\|h\|_{\partial^{-1}(\partial\mathcal{D} \odot \mathcal{D})} = \inf \left\{ \sum \|f_j\|_{\mathcal{D}} \|g_j\|_{\mathcal{D}} : h' = \sum f'_j g_j \right\}.$$

We next introduce the space \mathcal{X} which plays a role in the Dirichlet space theory analogous to the role of BMO in the Hardy space theory. We say $f \in \mathcal{X}$ if

$$\|f\|_{\mathcal{X}}^2 = |f(0)|^2 + \left\| |f'|^2 dA \right\|_{CM(\mathcal{D})} < \infty.$$

We denote the closure in \mathcal{X} of the space of polynomials by \mathcal{X}_0 .

Here is a summary of relations between the spaces. The duality pairings are with respect to the Dirichlet pairing $\langle \cdot, \cdot \rangle_{\mathcal{D}}$.

Theorem 23. *We have*

- (1) $\mathcal{X}_0^* = \mathcal{D} \odot \mathcal{D}$,
- (2) $(\mathcal{D} \odot \mathcal{D})^* = \mathcal{X}$,
- (3) $\mathcal{M}(\mathcal{D}) = H^\infty \cap \mathcal{X}$,
- (4) $\mathcal{D} \odot \mathcal{D} = \partial^{-1}(\partial\mathcal{D} \odot \mathcal{D})$.

Proof discussion. As we mentioned (3) is proved in [50].

A result essentially equivalent to $(\partial^{-1}(\partial\mathcal{D} \odot \mathcal{D}))^* = \mathcal{X}$ was proved by Coifman-Muri [30] using real variable techniques and in more function theoretic contexts by Tolokonnikov [53] and by Rochberg-Wu [43]. An interesting alternative approach to the result is given by Treil and Volberg in [54].

In [60] it is shown that $\mathcal{X}_0^* = \partial^{-1}(\partial\mathcal{D} \odot \mathcal{D})$. Item (2) is proved in [16] and when that is combined with the other results we obtain (1) and (4). \square

Statement (2) of the theorem is the analog of Nehari's characterization of bounded Hankel forms on the Hardy space, recast using the identification $H^2 \odot H^2 = H^1$ and Fefferman's duality theorem. Item (1) is the analog of Hartman's characterization of compact Hankel forms. Statement (4) is similar in spirit to the weak factorization result for Hardy spaces given by Aleksandrov and Peller in [4] where they study Foguel-Hankel operators on the Hardy space.

Given the previous theorem it is easy to check the inclusions

$$(40) \quad \mathcal{M}(\mathcal{D}) \subset \mathcal{X} \subset \mathcal{D} \subset \mathcal{D} \odot \mathcal{D}$$

In our paper [17] we discuss more facts about these spaces.

6.3. The Corona Theorem. In 1962 Lennart Carleson demonstrated in [26] the absence of a corona in the maximal ideal space of $H^\infty(\mathbb{D})$ by showing that if $\{g_j\}_{j=1}^N$ is a finite set of functions in $H^\infty(\mathbb{D})$ satisfying

$$(41) \quad \sum_{j=1}^N |g_j(z)| \geq \delta > 0, \quad z \in \mathbb{D},$$

then there are functions $\{f_j\}_{j=1}^N$ in $H^\infty(\mathbb{D})$ with

$$(42) \quad \sum_{j=1}^N f_j(z) g_j(z) = 1, \quad z \in \mathbb{D},$$

While not immediately obvious, the result of Carleson is in fact equivalent to the following statement about the Hilbert space $H^2(\mathbb{D})$. If one is given a finite set of functions $\{g_j\}_{j=1}^N$ in $H^\infty(\mathbb{D})$ satisfying (41) and a function $h \in H^2(\mathbb{D})$, then there are functions $\{f_j\}_{j=1}^N$ in $H^2(\mathbb{D})$ with

$$(43) \quad \sum_{j=1}^N f_j(z) g_j(z) = h(z), \quad z \in \mathbb{D},$$

The key difference between (42) and (43) is that one is solving the problem in the Hilbert space setting as opposed to the multiplier algebra, which makes the problem somewhat easier.

In this section we discuss the Corona Theorem for the multiplier algebra of the Dirichlet space. The method of proof will be intimately connected with the resulting statements for $H^\infty(\mathbb{D})$ and $H^2(\mathbb{D})$. We also will connect this result to a related statement for the Hilbert space \mathcal{D} . The proof of this fact is given by $\bar{\partial}$ -methods and the connections between weak Carleson measures for the space $\mathcal{D}(\mathbb{D})$. Another proof can be given by simply proving the Hilbert space version directly and then applying the Toeplitz Corona Theorem. Implicit in both versions are certain solutions to $\bar{\partial}$ -problems that arise.

6.3.1. *The $\bar{\partial}$ -equation in the Dirichlet Space.* As is well-known there is an intimate connection between the Corona Theorem and $\bar{\partial}$ -problems. In our context, a $\bar{\partial}$ -problem will be to solve the following differential equation

$$(44) \quad \bar{\partial}b = \mu$$

where μ is a Carleson measure for the space \mathcal{D} and b is some unknown function. Now solving this problem is an easy application of Cauchy's formula, however we will need to obtain estimates of the solutions. To obtain these estimates, one needs a different solution operator to the $\bar{\partial}$ -problem more appropriately suited to our contexts.

In [61] Xiao's constructed a non-linear solution operator for (44) that is well adapted to solve (44) and obtain estimates. We note that in the case of $H^\infty(\mathbb{D})$ that this result was first obtained by P. Jones, [33]. First, note that

$$F(z) = \frac{1}{2\pi i} \iint_{\mathbb{D}} \frac{d\mu(\zeta)}{\zeta - z} d\zeta \wedge d\bar{\zeta}$$

satisfies $\bar{\partial}F = \mu$ in the sense of distribution.

The difficulty with this solution kernel is that it does not allow for one to obtain good estimates on the solution. To rectify this, following Jones [33], we define a new non-linear kernel that will overcome this difficulty.

Theorem 24 (Jones, [33]). *Let μ be a complex $H^2(\mathbb{D})$ Carleson measure on \mathbb{D} . Then with $S(\mu)(z)$ given by*

$$(45) \quad S(\mu)(z) = \iint_{\mathbb{D}} K(\sigma, z, \zeta) d\nu(\zeta)$$

where $\sigma = \frac{|\mu|}{\|\mu\|_{CM(H^2)}}$ and

$$K(\sigma, z, \zeta) \equiv \frac{2i}{\pi} \frac{1 - |\zeta|^2}{(z - \zeta)(1 - \bar{\zeta}z)} \exp \left\{ \iint_{|\omega| \geq |\zeta|} \left(-\frac{1 + \bar{\omega}z}{1 - \bar{\omega}z} + \frac{1 + \bar{\omega}\zeta}{1 - \bar{\omega}\zeta} \right) d\sigma(\omega) \right\},$$

we have that:

- (1) $S(\mu) \in L^1_{loc}(\mathbb{D})$.
- (2) $\bar{\partial}S(\mu) = \mu$ in the sense of distributions.
- (3) $\iint_{\mathbb{D}} \left| K\left(\frac{|\mu|}{\|\mu\|_{C^{\alpha r}}}, x, \zeta\right) \right| d|\mu|(\zeta) \lesssim \|\mu\|_{CM(H^2)}$ for all $x \in \mathbb{T} = \partial\mathbb{D}$,

$$\text{so } \|S(\mu)\|_{L^\infty(\mathbb{T})} \lesssim \|\mu\|_{CM(H^2)}.$$

With this set-up, we now state the follow theorem due to Xiao, extending Theorem 24, about estimates for $\bar{\partial}$ -problems in the Dirichlet space.

Theorem 25 (Xiao, [61]). *If $|g(z)|^2 dA(z)$ is a \mathcal{D} -Carleson measure then the function $S(\mu)(z)$ satisfies $\bar{\partial}S(\mu) = \mu$ and*

$$\|S(\mu)\|_{\mathcal{W}^{1/2}(\mathbb{T})} \lesssim \|\mu\|_{CM(\mathcal{D})}$$

6.3.2. Corona Theorems and Complete Nevanlinna-Pick Kernels. Let X be a Hilbert space of holomorphic functions in an open set Ω in \mathbb{C}^n that is a reproducing kernel Hilbert space with a *complete irreducible Nevanlinna-Pick kernel* (see [2] for the definition). The following *Toeplitz corona theorem* is due to Ball, Trent and Vinnikov [19] (see also Ambrozie and Timotin [5] and Theorem 8.57 in [2]).

For $f = (f_\alpha)_{\alpha=1}^N \in \oplus^N X$ and $h \in X$, define $\mathbb{M}_f h = (f_\alpha h)_{\alpha=1}^N$ and

$$\|f\|_{Mult(X, \oplus^N X)} = \|\mathbb{M}_f\|_{X \rightarrow \oplus^N X} = \sup_{\|h\|_X \leq 1} \|\mathbb{M}_f h\|_{\oplus^N X}.$$

Note that $\max_{1 \leq \alpha \leq N} \|\mathcal{M}_{f_\alpha}\|_{M_X} \leq \|f\|_{Mult(X, \oplus^N X)} \leq \sqrt{\sum_{\alpha=1}^N \|\mathcal{M}_{f_\alpha}\|_{M_X}^2}$.

Theorem 26 (Toeplitz Corona Theorem). *Let X be a Hilbert function space in an open set Ω in \mathbb{C}^n with an irreducible complete Nevanlinna-Pick kernel. Let $\delta > 0$ and $N \in \mathbb{N}$. Then $g_1, \dots, g_N \in M_X$ satisfy the following “baby corona property”; for every $h \in X$, there are $f_1, \dots, f_N \in X$ such that*

$$(46) \quad \begin{aligned} \|f_1\|_X^2 + \dots + \|f_N\|_X^2 &\leq \frac{1}{\delta} \|h\|_X^2, \\ g_1(z) f_1(z) + \dots + g_N(z) f_N(z) &= h(z), \quad z \in \Omega, \end{aligned}$$

if and only if $g_1, \dots, g_N \in M_X$ satisfy the following “multiplier corona property”; there are $\varphi_1, \dots, \varphi_N \in M_X$ such that

$$(47) \quad \begin{aligned} \|\varphi\|_{Mult(X, \oplus^N X)} &\leq 1, \\ g_1(z) \varphi_1(z) + \dots + g_N(z) \varphi_N(z) &= \sqrt{\delta}, \quad z \in \Omega. \end{aligned}$$

The *baby corona theorem* is said to hold for X if whenever $g_1, \dots, g_N \in M_X$ satisfy

$$(48) \quad |g_1(z)|^2 + \dots + |g_N(z)|^2 \geq c > 0, \quad z \in \Omega,$$

then g_1, \dots, g_N satisfy the baby corona property (46).

We now state a simple proposition that will be useful in understanding the relationships between the Corona problems for \mathcal{D} and $M_{\mathcal{D}}$.

Proposition 27. *Suppose that $g_1, \dots, g_N \in M_{\mathcal{D}}$. Define the map*

$$M_{(g_1, \dots, g_N)}(f_1, \dots, f_N) := \sum_{k=1}^N g_k(z) f_k(z).$$

Then the following are equivalent

- (i) $M_{(g_1, \dots, g_n)} : M_{\mathcal{D}} \times \cdots \times M_{\mathcal{D}} \mapsto M_{\mathcal{D}}$ is onto;
- (ii) $M_{(g_1, \dots, g_n)} : \mathcal{D} \times \cdots \times \mathcal{D} \mapsto \mathcal{D}$ is onto;
- (iii) There exists a $\delta > 0$ such that for all $z \in \mathbb{D}$ we have

$$\sum_{k=1}^N |g_k(z)|^2 \geq \delta > 0.$$

It is easy to see that both (i) and (ii) each individually imply (iii). We will show that condition (iii) implies both (i) and (ii). Note that by the Toeplitz Corona Theorem 26 it would suffice to prove that (iii) implies (ii) since the result then lifts to give the statement in (i). The proof of Proposition 27 follows by the lines of Wolff's proof of the Corona Theorem for $H^\infty(\mathbb{D})$, but uses the solution operator given by Xiao in Theorem 25.

It is important to point out that there are several other proof of Proposition 27 at this point. The first proof of this fact was given by Tolokonnikov, [53] and was essentially obtained via connections with Carleson's Corona Theorem. Another proof of this result, but with the added benefit of being true for an infinite number of generators was given by Trent [55]. Trent demonstrated that (iii) implies (ii), and then applied the Toeplitz Corona Theorem to deduce that (iii) implies (i). This proof exploits the fact that the kernel for the Dirichlet space is a complete Nevanlinna-Pick kernel. Finally, there is a more recent proof of this fact by Costea, Sawyer and Wick, [31]. The method of proof again is to demonstrate the Corona Theorem for \mathcal{D} under the hypothesis (iii). The proof in [31] is true more generally for the Dirichlet space in any dimension.

7. INTERPOLATING SEQUENCES

Let \mathcal{H} be a reproducing kernel Hilbert space (RKHS) of functions defined on some space X , with kernel functions $\{k_z\}_{z \in X}$. Let $\mathcal{M}(\mathcal{H})$ be the multiplier space of \mathcal{H} . A sequence $S \subseteq X$ is an *interpolating sequence for the multiplier algebra* $\mathcal{M}(\mathcal{H})$ if the restriction map

$$R_S : g \mapsto \{g(s) : s \in S\}$$

maps $\mathcal{M}(\mathcal{H})$ onto ℓ^∞ . Since $\mathcal{M}(\mathcal{H}) \subseteq L^\infty(X)$, the map is automatically bounded. Consider the weight $w : S \rightarrow \mathbb{R}^+$, $w(s) = \|k_s\|_{\mathcal{H}}^{-2}$. We say that the sequence S is an *interpolating sequence for the space* \mathcal{H} if R_S is a bounded map of \mathcal{H} onto $\ell^2(S, w)$. In the context of complete Nevanlinna-Pick RKHS these two notions coincide [37]. [Our terminology differs from some sources. Bishop [21], for instance, calls *universally interpolating sequences* for \mathcal{D} what we call interpolating sequences, and simply calls *interpolating sequences what we will call onto interpolating sequences.*]

Theorem 28 (Marshall-Sundberg, [37]). *Let \mathcal{H} be a RKHS of functions on some space X , with the complete Nevanlinna-Pick property. For a sequence S the following are equivalent:*

- (1) S is interpolating for $\mathcal{M}(\mathcal{H})$.
- (2) S is interpolating for \mathcal{H} .
- (3) The family of functions $\left\{ \frac{k_s}{\|k_s\|_{\mathcal{H}}} \right\}_{s \in S}$ is a **Riesz basis** for the space \mathcal{H} :

$$\left\| \sum_{s \in S} a_s \frac{k_s}{\|k_s\|_{\mathcal{H}}} \right\|_{\mathcal{H}}^2 \approx \sum_{s \in S} |a_s|^2.$$

Sarason observed that interpolating sequences for the multiplier space $\mathcal{M}(\mathcal{H})$ have a distinguished rôle in the theory of the RKHS space \mathcal{H} . Let φ be a multiplier of the space \mathcal{H} and $S = \{s_j : j = 1, \dots, n\}$ be a sequence in X . Let M_φ the multiplication operator by φ and M_φ^* be its adjoint. Then, $\{\overline{\varphi(s_j)} : j = 1, \dots, n\}$ is a set of eigenvalues for M_φ^* , having the corresponding kernel functions as eigenvectors: $M_\varphi^* k_{s_j} = \overline{\varphi(s_j)} k_{s_j}$.

Finding the multiplier φ which interpolates data $\varphi(s_j) = \lambda_j$ corresponds, then, to extending the diagonal operator $k_{s_j} \mapsto \overline{\lambda_j} k_{s_j}$ (which is defined on $\text{span}\{k_{s_j} : j=1, \dots, n\}$) to the adjoint of a multiplication operator. We redirect the interested reader to the book [2], to the article [3] and to the important manuscript [37] itself for far reaching developments of this line of reasoning.

For a given sequence S in X , there are two obvious *necessary conditions* for it to be interpolating for \mathcal{H} :

- (Sep) The sequence S is **separated**: there is a positive $\sigma < 1$ such that for all $s, t \in S$ one has

$$\left| \left\langle \frac{k_s}{\|k_s\|_{\mathcal{H}}}, \frac{k_t}{\|k_t\|_{\mathcal{H}}} \right\rangle \right| \leq \sigma$$

This condition expresses the fact that there exists a function $f \in \mathcal{H}$ such that $f(s) = 0$ and $f(t) = 1$.

- (CM) The measure $\mu_S = \sum_{s \in S} \|k_s\|_{\mathcal{H}}^{-2} \delta_s$ is a **Carleson measure for the space \mathcal{H}** :

$$\int_X |f|^2 d\mu_S \leq C(\mu) \|f\|_{\mathcal{H}}^2,$$

which expresses the boundedness of the restriction map R_S .

Kristian Seip [47] conjectures that, for RKHS with the complete Nevanlinna-Pick property, these two conditions are sufficient for S to be interpolating. Carleson's celebrated Interpolation [28] Theorem says that such is the case when $\mathcal{H} = H^2$ is the Hardy space. Bøe proved Seip's conjecture under an additional assumption on the kernel functions (an assumption which, interestingly, is *not* satisfied by the Hardy space itself, but which is satisfied by the Dirichlet space).

7.1. Interpolating Sequences for \mathcal{D} and its Multiplier Space. The characterization of the interpolating functions was independently solved by Marshall and Sundberg [37] and by Bishop [21] in 1994.

Theorem 29 (Marshall-Sundberg [37], Bishop [21]). *A sequence S in \mathbb{D} is interpolating for the Dirichlet space \mathcal{D} if and only if it satisfies (Sep) and (CM).*

Actually, Bishop proved that interpolating sequences for \mathcal{D} are also interpolating for $\mathcal{M}(\mathcal{D})$, but not the converse. At the present moment, there are four essentially different proofs that (Sep) and (CM) are necessary and sufficient for \mathcal{D} interpolation: [37], [21], [25], [24]. Interpolating sequences for the Dirichlet space differ in one important aspect from interpolating sequences for the Hardy space. In the case of H^2 , in fact, if the restriction operator is surjective (if, in our terminology, the sequence S is onto interpolating), then it is automatically bounded. As we will see in the next subsection, there are sequences S in the unit disc for which the restriction operator is surjective, but not bounded.

It is interesting and useful to restate the separation condition in terms of hyperbolic distance: *(Sep) in the Dirichlet space \mathcal{D} holds for the sequence S in \mathbb{D} if and*

only if there are positive constants A, B such that, for all $z \neq w \in S$

$$\max\{d(z), d(w)\} \leq Ad(z, w) + B.$$

This huge separation, which is related to the hyperbolic invariance of the Dirichlet norm, compensates -in the solution of the interpolating sequences problem and in other questions- for the lack of Blaschke products. In fact, it allows much space for crafting holomorphic functions from smooth ones, with little overlapping.

Proof(s) Discussion. [37]. In their article, Marshall and Sundberg first developed a general theory concerning interpolating sequences in spaces with the complete Nevanlinna-Pick property. In particular, they reduced the problem of characterizing the interpolating sequences for \mathcal{D} to that of the interpolating sequences for its multiplier space. This left them with the (hard) task of showing that, given (Sep) and (CM), one could interpolate bounded sequences by multiplier functions. In order to do that, they first solved the easier (but still difficult) problem of interpolating the data by means of a smooth function $\varphi : \mathbb{D} \rightarrow \mathbb{R}$, having properties similar to those of a multiplier in $\mathcal{M}(\mathcal{D})$: φ is bounded, it has finite Dirichlet norm and, more, $|\nabla\varphi|^2 dA \in CM(\mathcal{D})$.

The basic building block for constructing such φ are functions φ_z attached to points z in \mathbb{D} , which are, substantially, the best smoothed version (in terms of Dirichlet integral) of the function $\chi_{\tilde{S}(z)}$, where

$$\tilde{S}(z) = \left\{ w \in \mathbb{D} : \left| w - \frac{z}{|z|} \right| \leq (1 - |z|)^\alpha \right\}$$

($\alpha < 1$ suitably chosen) is the “enlarged Carleson box” having center in z . The separation condition (Sep) ensures that, if z_1, z_2 are points of the sequence S and $\text{supp}(\varphi_{z_1}) \cap \text{supp}(\varphi_{z_2}) \neq \emptyset$, then one of the points has to be much closer to the boundary than the other. This is one of the two main tools (the other being the Carleson measure condition, which further separates the points of the sequence) in the various estimates for linear combinations of basic building functions. These basic building blocks and their holomorphic modifications are the main tool in the proofs of the interpolating theorems in [25] and [12].

The rest of the proof consists in showing that one can correct the function φ , making it harmonic, and from this one easily proceeds to the holomorphic case.

Bishop, instead, uses as building blocks conformal maps, (see [21], p.27). In his article, he observes that the construction of the interpolating functions for \mathcal{D} does not require the full use of the assumption (CM): contrary to the Hardy case, there are sequences S for which the restriction operator is *onto and unbounded*. We will return on this in the next section.

Böe’s short proof in [24] is less constructive, and it relies on Hilbert space arguments. However, in his paper [25], dealing with the more general case of the analytic Besov space, Böe has an explicit construction of the interpolating sequences for \mathcal{D} and $\mathcal{M}(\mathcal{D})$. He makes use of holomorphic modifications of the functions φ_z in [37], which are the starting point for a hard and clever recursion scheme.

It is clear from the construction in [24] that, under the assumptions (Sep) and (CM), one has *linear interpolation of data*, both in \mathcal{D} and $\mathcal{M}(\mathcal{D})$: there exist linear operators $L_S : \ell^\infty(S) \rightarrow \mathcal{M}(\mathcal{D})$ and $\Lambda_S : \ell^2(S, w) \rightarrow \mathcal{D}$ such that $L_S\{a_s : s \in S\}$ solves the interpolating problem in $\mathcal{M}(\mathcal{D})$ with data $\{a_s : s \in S\} \in \ell^\infty(S)$ and

$\Lambda_S\{b_s : s \in S\}$ solves the interpolating problem in \mathcal{D} with data $\{b_s : s \in S\} \in \ell^2(S, w)$. \square

7.2. Weak Interpolation and “Onto” Interpolation. A sequence S in \mathbb{D} is *onto interpolating* if the restriction operator R_S maps \mathcal{D} onto $\ell^2(S, w)$. We do not ask the operator R_S to be bounded (hence, to be defined on all of \mathcal{D}). It follows from the Closed Graph Theorem that, if S is onto interpolating, then it is interpolating *with norm control*: there is a constant $C > 0$ such that for $\{a_s : s \in S\} \in \ell^2(S, w)$ there is $f \in \mathcal{D}$ such that $f(s) = a_s$ and $\|f\|_{\mathcal{D}} \leq C\|\{a_s : s \in S\}\|_{\ell^2(S, w)}$. Furthermore, interpolation can be realized linearly.

A sequence S in \mathbb{D} is *weakly interpolating* if there is $C > 0$ such that, for all $s_0 \in S$ there is $f_{s_0} \in \mathcal{D}$ with $f_{s_0}(s) = \delta_{s_0}(s)$ for $s \in S$ (δ_{s_0} is the Kroenecker function) and norm control $\|f_{s_0}\|_{\mathcal{D}}^2 \leq C\|\delta_{s_0}\|_{\ell^2(S, w)}^2 \approx d(s_0)^{-1}$. Clearly, weakly interpolating is weaker than onto interpolating.

Remark 30. (a) *Weak interpolation (a fortiori, onto interpolation) implies the separation condition (Sep).*

(b) *By adding a finite number of points to an onto interpolating sequence, we obtain another onto interpolating sequence.*

A geometric characterization of the onto interpolating sequences is still lacking. However, the following facts are known.

Theorem 31 (Bishop [21]). *The sequence S is onto interpolating if and only if it is weakly interpolating, and this is in turn equivalent to having weak interpolation with functions which satisfy the further condition that $\|f_s\|_{L^\infty(\mathcal{D})} \leq C$ for some constant C .*

The proof of Bishop’s Theorem involves the clever use of a variety of sophisticated tools. It would be interesting having a different proof (one which worked for the analytic Besov spaces, for instance). Unfortunately, establishing whether a given sequence S is weakly interpolating is not much easier than establishing if it is onto interpolating.

Both Bishop [21] and Bøe realized that a sequence S is onto interpolating if the associated measure μ_S satisfies the *simple condition* (13) instead of the stronger Carleson measure condition (CM). The simple condition implies, in particular, that the measure μ_S is finite and Bishop asked whether there are onto interpolating sequences with infinite μ_S . The answer is affirmative:

Theorem 32 (Arcozzi-Rochberg-Sawyer, [12]). *There exist sequences S in \mathbb{D} with $\mu_S(\mathbb{D}) = +\infty$, which are onto interpolating for \mathcal{D} .*

The proof of Theorem 32 relies on a modification of Bøe’s recursive scheme, using Bøe’s functions.

In [12] there is another partial result, which extends the theorems of Bishop and Bøe. In order to state it, we have to go back to the tree language. Let \mathcal{T} be the dyadic tree associated with the disc \mathbb{D} . By Remarks 30, we can assume that each box α in \mathcal{T} contains at most one point from the candidate interpolating sequence S in \mathbb{D} . We can therefore identify points in S with distinguished boxes in \mathcal{T} . We say that S in \mathbb{D} satisfies the *weak simple condition* if for all α in \mathcal{T} ,

$$(49) \quad \sum_{\substack{\beta \in S, \beta > \alpha \\ \mu_S(\gamma) = 0 \text{ for } \alpha < \gamma < \beta}} \mu_S(\beta) \leq Cd(\alpha)^{-1}.$$

Theorem 33 ([12]). *Let S be a sequence in \mathbb{D} and suppose that $\mu_S(\mathbb{D}) < \infty$. If μ_S satisfies (Sep) and the weak simple condition (49), then S is onto interpolating for \mathcal{D} .*

We observe that the weak simple condition can not be necessary for onto interpolation. In fact, it is easy to produce examples of sequences S satisfying (49), having subsequences S' (which are then onto interpolating for \mathcal{D}) for which (49) is not satisfied. Such examples, however, have $\mu_S(\mathbb{D}) = +\infty$. We do not know whether, under the assumptions $\mu_S(\mathbb{D}) = +\infty$ and (Sep), the weak simple condition is necessary for onto interpolation.

As a last, partial result about onto interpolation, let us mention a necessary condition of capacity type. If S an onto interpolating sequence for \mathcal{D} in \mathbb{D} , which we might identify with a subsequence of the tree \mathcal{T} , then the *discrete capacity condition holds*:

to each s_0 in S , there corresponds a positive function φ_{s_0} on \mathcal{T} such that $\|\varphi_{s_0}\|_{\ell^2}^2 \leq Cd(s_0)^{-1}$ and $\varphi_{s_0}(s) = \delta_{s_0}(s)$ whenever $s \in S$.

A proof of this fact easily follows from Proposition 8. The discrete capacity condition can be stated in terms of discrete condenser capacities:

$$\text{Cap}_{\mathcal{T}}(S \setminus \{s_0\}, \{s_0\}) \leq Cd(s_0)^{-1}.$$

A reasonable conjecture is that the discrete capacity condition, plus the separation condition (Sep), are necessary and sufficient in order for S to be onto interpolating. Other material on the problem of the onto interpolating sequences is in [12].

REFERENCES

- [1] David R. Adams, *On the existence of capacity strong type estimates in R^n* , Ark. Mat. **14** (1976), no. 1, 125–140.
- [2] Jim Agler and John E. McCarthy, *Pick interpolation and Hilbert function spaces*, Graduate Studies in Mathematics, vol. 44, American Mathematical Society, Providence, RI, 2002.
- [3] ———, *Complete Nevanlinna-Pick kernels*, J. Funct. Anal. **175** (2000), no. 1, 111–124.
- [4] A. B. Aleksandrov and V. V. Peller, *Hankel operators and similarity to a contraction*, Internat. Math. Res. Notices (1996), no. 6, 263–275.
- [5] Călin-Grigore Ambrozie and Dan Timotin, *On an intertwining lifting theorem for certain reproducing kernel Hilbert spaces*, Integral Equations Operator Theory **42** (2002), no. 4, 373–384.
- [6] J. Arazy and S. D. Fisher, *The uniqueness of the Dirichlet space among Möbius-invariant Hilbert spaces*, Illinois J. Math. **29** (1985), no. 3, 449–462.
- [7] J. Arazy, S. D. Fisher, and J. Peetre, *Möbius invariant function spaces*, J. Reine Angew. Math. **363** (1985), 110–145.
- [8] Nicola Arcozzi, *Carleson measures for analytic Besov spaces: the upper triangle case*, JIPAM. J. Inequal. Pure Appl. Math. **6** (2005), no. 1, Article 13, 15 pp. (electronic).
- [9] Nicola Arcozzi and Richard Rochberg, *Topics in dyadic Dirichlet spaces*, New York J. Math. **10** (2004), 45–67 (electronic).

- [10] N. Arcozzi, R. Rochberg, and E. Sawyer, *Carleson measures for the Drury-Arveson Hardy space and other Besov-Sobolev spaces on complex balls*, Adv. Math. **218** (2008), no. 4, 1107–1180.
- [11] ———, *Capacity, Carleson measures, boundary convergence, and exceptional sets*, Perspectives in partial differential equations, harmonic analysis and applications, Proc. Sympos. Pure Math., vol. 79, Amer. Math. Soc., Providence, RI, 2008, pp. 1–20.
- [12] ———, *Onto interpolating sequences for the Dirichlet space*, preprint pp., available at <http://amsacta.cib.unibo.it/2480/>.
- [13] Nicola Arcozzi, Richard Rochberg, and Eric Sawyer, *The characterization of the Carleson measures for analytic Besov spaces: a simple proof*, Complex and harmonic analysis, DEStech Publ., Inc., Lancaster, PA, 2007, pp. 167–177.
- [14] N. Arcozzi, R. Rochberg, and E. Sawyer, *Carleson measures and interpolating sequences for Besov spaces on complex balls*, Mem. Amer. Math. Soc. **182** (2006), no. 859, vi+163.
- [15] Nicola Arcozzi, Richard Rochberg, and Eric Sawyer, *Carleson measures for analytic Besov spaces*, Rev. Mat. Iberoamericana **18** (2002), no. 2, 443–510.
- [16] Nicola Arcozzi, Richard Rochberg, Eric Sawyer, and Brett D. Wick, *Bilinear forms on the Dirichlet space*, Anal. & PDE **3** (2010), no. 1, 21–47.
- [17] ———, *Function spaces related to the Dirichlet space*, J. London Math. Soc., available at <http://arxiv.org/abs/0812.3422>.
- [18] ———, *Potential Theory on Trees, Graphs and Ahlfors regular metric spaces*, preprint.
- [19] Joseph A. Ball, Tavan T. Trent, and Victor Vinnikov, *Interpolation and commutant lifting for multipliers on reproducing kernel Hilbert spaces*, Operator theory and analysis (Amsterdam, 1997), Oper. Theory Adv. Appl., vol. 122, Birkhäuser, Basel, 2001, pp. 89–138.
- [20] Arne Beurling, *Ensembles exceptionnels*, Acta Math. **72** (1940), 1–13 (French).
- [21] C. J. Beurling, *Interpolating sequences for the Dirichlet space and its multipliers*, preprint pp., available at <http://www.math.sunysb.edu/~bishop/papers/mult.pdf>.
- [22] Itai Benjamini and Yuval Peres, *A correlation inequality for tree-indexed Markov chains*, Seminar on Stochastic Processes, 1991 (Los Angeles, CA, 1991), Progr. Probab., vol. 29, Birkhäuser Boston, Boston, MA, 1992, pp. 7–14.
- [23] Daniel Blasi and Jordi Pau, *A characterization of Besov-type spaces and applications to Hankel-type operators*, Michigan Math. J. **56** (2008), no. 2, 401–417.
- [24] Bjarte Bøe, *A norm on the holomorphic Besov space*, Proc. Amer. Math. Soc. **131** (2003), no. 1, 235–241 (electronic).
- [25] ———, *Interpolating sequences for Besov spaces*, J. Funct. Anal. **192** (2002), no. 2, 319–341.
- [26] Lennart Carleson, *Interpolations by bounded analytic functions and the corona problem*, Ann. of Math. (2) **76** (1962), 547–559.
- [27] ———, *A representation formula for the Dirichlet integral*, Math. Z. **73** (1960), 190–196.
- [28] ———, *An interpolation problem for bounded analytic functions*, Amer. J. Math. **80** (1958), 921–930.
- [29] ———, *On the zeros of functions with bounded Dirichlet integrals*, Math. Z. **56** (1952), 289–295.

- [30] Ronald R. Coifman and Takafumi Murai, *Commutators on the potential-theoretic energy spaces*, Tohoku Math. J. (2) **40** (1988), no. 3, 397–407.
- [31] Şerban Costea, Eric T. Sawyer, and Brett D. Wick, *The Corona Theorem for the Drury-Arveson Hardy space and other holomorphic Besov-Sobolev spaces on the unit ball in \mathbb{C}^n* , Anal. & PDE, to appear, available at <http://arxiv.org/abs/0811.0627>.
- [32] L. I. Hedberg and Th. H. Wolff, *Thin sets in nonlinear potential theory*, Ann. Inst. Fourier (Grenoble) **33** (1983), no. 4, 161–187.
- [33] Peter W. Jones, *L^∞ estimates for the $\bar{\partial}$ problem in a half-plane*, Acta Math. **150** (1983), no. 1-2, 137–152.
- [34] N. J. Kalton and I. E. Verbitsky, *Nonlinear equations and weighted norm inequalities*, Trans. Amer. Math. Soc. **351** (1999), no. 9, 3441–3497.
- [35] Ron Kerman and Eric Sawyer, *Carleson measures and multipliers of Dirichlet-type spaces*, Trans. Amer. Math. Soc. **309** (1988), no. 1, 87–98.
- [36] R. Kerman and E. Sawyer, *The trace inequality and eigenvalue estimates for Schrödinger operators*, Ann. Inst. Fourier (Grenoble) **36** (1986), no. 4, 207–228 (English, with French summary).
- [37] D. Marshall and C. Sundberg, *Interpolating sequences for the multipliers of the Dirichlet space*, preprint pp., available at <http://www.math.washington.edu/~marshall/preprints/interp.pdf>.
- [38] V. G. Maz'ja, *Certain integral inequalities for functions of several variables*, Problems of mathematical analysis, No. 3: Integral and differential operators, Differential equations (Russian), Izdat. Leningrad. Univ., Leningrad, 1972, pp. 33–68 (Russian).
- [39] Benjamin Muckenhoupt and Richard Wheeden, *Weighted norm inequalities for fractional integrals*, Trans. Amer. Math. Soc. **192** (1974), 261–274.
- [40] Subhashis Nag and Dennis Sullivan, *Teichmüller theory and the universal period mapping via quantum calculus and the $H^{1/2}$ space on the circle*, Osaka J. Math. **32** (1995), no. 1, 1–34.
- [41] Alexander Nagel, Walter Rudin, and Joel H. Shapiro, *Tangential boundary behavior of functions in Dirichlet-type spaces*, Ann. of Math. (2) **116** (1982), no. 2, 331–360.
- [42] Stefan Richter and Carl Sundberg, *A formula for the local Dirichlet integral*, Michigan Math. J. **38** (1991), no. 3, 355–379.
- [43] Richard Rochberg and Zhi Jian Wu, *A new characterization of Dirichlet type spaces and applications*, Illinois J. Math. **37** (1993), no. 1, 101–122.
- [44] William T. Ross, *The classical Dirichlet space*, Recent advances in operator-related function theory, Contemp. Math., vol. 393, Amer. Math. Soc., Providence, RI, 2006, pp. 171–197.
- [45] Donald Sarason, *Holomorphic spaces: a brief and selective survey*, Holomorphic spaces (Berkeley, CA, 1995), Math. Sci. Res. Inst. Publ., vol. 33, Cambridge Univ. Press, Cambridge, 1998, pp. 1–34.
- [46] Stanley A. Sawyer, *Martin boundaries and random walks*, Harmonic functions on trees and buildings (New York, 1995), Contemp. Math., vol. 206, Amer. Math. Soc., Providence, RI, 1997, pp. 17–44.
- [47] Kristian Seip, *Interpolation and sampling in spaces of analytic functions*, University Lecture Series, vol. 33, American Mathematical Society, Providence, RI, 2004.

- [48] H. S. Shapiro and A. L. Shields, *On the zeros of functions with finite Dirichlet integral and some related function spaces*, Math. Z. **80** (1962), 217–229.
- [49] ———, *On some interpolation problems for analytic functions*, Amer. J. Math. **83** (1961), 513–532.
- [50] David A. Stegenga, *Multipliers of the Dirichlet space*, Illinois J. Math. **24** (1980), no. 1, 113–139.
- [51] Elias M. Stein, *Singular integrals and differentiability properties of functions*, Princeton Mathematical Series, No. 30, Princeton University Press, Princeton, N.J., 1970.
- [52] Edgar Tchoundja, *Carleson measures for the generalized Bergman spaces via a $T(1)$ -type theorem*, Ark. Mat. **46** (2008), no. 2, 377–406.
- [53] V. A. Tolokonnikov, *Carleson’s Blaschke products and Douglas algebras*, Algebra i Analiz **3** (1991), no. 4, 186–197 (Russian); English transl., St. Petersburg Math. J. **3** (1992), no. 4, 881–892.
- [54] Sergei Treil and Alexander Volberg, *A fixed point approach to Nehari’s problem and its applications*, Toeplitz operators and related topics (Santa Cruz, CA, 1992), Oper. Theory Adv. Appl., vol. 71, Birkhäuser, Basel, 1994, pp. 165–186.
- [55] Tavan T. Trent, *A corona theorem for multipliers on Dirichlet space*, Integral Equations Operator Theory **49** (2004), no. 1, 123–139.
- [56] J. B. Twomey, *Tangential boundary behaviour of harmonic and holomorphic functions*, J. London Math. Soc. (2) **65** (2002), no. 1, 68–84.
- [57] I. È. Verbitskiĭ, *Multipliers in spaces with “fractional” norms, and inner functions*, Sibirsk. Mat. Zh. **26** (1985), no. 2, 51–72, 221 (Russian).
- [58] Alexander Volberg and Brett D. Wick, *Bergman-type Singular Operators and the Characterization of Carleson Measures for Besov–Sobolev Spaces on the Complex Ball*, Amer. J. Math., to appear, available at <http://arxiv.org/abs/0910.1142>.
- [59] Zhijian Wu, *Function theory and operator theory on the Dirichlet space*, Holomorphic spaces (Berkeley, CA, 1995), Math. Sci. Res. Inst. Publ., vol. 33, Cambridge Univ. Press, Cambridge, 1998, pp. 179–199.
- [60] ———, *The predual and second predual of W_α* , J. Funct. Anal. **116** (1993), no. 2, 314–334.
- [61] Jie Xiao, *The $\bar{\partial}$ -problem for multipliers of the Sobolev space*, Manuscripta Math. **97** (1998), no. 2, 217–232.
- [62] Kehe Zhu, *Operator theory in function spaces*, 2nd ed., Mathematical Surveys and Monographs, vol. 138, American Mathematical Society, Providence, RI, 2007.

N. ARCOZZI, DIPARTIMENTO DO MATEMATICA, UNIVERSITA DI BOLOGNA, 40127 BOLOGNA, ITALY

E-mail address: `arcozzi@dm.unibo.it`

R. ROCHBERG, DEPARTMENT OF MATHEMATICS, WASHINGTON UNIVERSITY, ST. LOUIS, MO 63130, U.S.A

E-mail address: `rr@math.wustl.edu`

E. T. SAWYER, DEPARTMENT OF MATHEMATICS & STATISTICS, MCMASTER UNIVERSITY, HAMILTON, ONTARIO, L8S 4K1, CANADA

E-mail address: `sawyer@mcmaster.ca`

B. D. WICK, SCHOOL OF MATHEMATICS, GEORGIAN INSTITUTE OF TECHNOLOGY, 686 CHERRY, STREET, ATLANTA, GA USA 30332-0160

E-mail address: `wick@math.gatech.edu`