

THE MINIMALLY DISPLACED SET OF AN IRREDUCIBLE AUTOMORPHISM OF F_N IS CO-COMPACT

STEFANO FRANCAVIGLIA

Dipartimento di Matematica of the University of Bologna

ARMANDO MARTINO

Mathematical Sciences, University of Southampton

DIONYSIOS SYRIGOS

Mathematical Sciences, University of Southampton

ABSTRACT. We study the minimally displaced set of irreducible automorphisms of a free group. Our main result is the co-compactness of the minimally displaced set of an irreducible automorphism with exponential growth ϕ , under the action of the centraliser $C(\phi)$. As a corollary, we get that the same holds for the action of $\langle \phi \rangle$ on $Min(\phi)$. Finally, we prove that the minimally displaced set of an irreducible automorphism of growth rate one is consisted of a single point.

CONTENTS

1. Introduction	1
2. Preliminaries	3
2.1. Culler -Vogtmann space	3
2.2. Stretching factor and Automorphisms	4
2.3. Properties of the Thick Part	6
2.4. Connectivity of the Min-set	6
3. Results	7
3.1. Exponential growth	7
3.2. Growth rate one	9
References	12

1. INTRODUCTION

In this paper, we study the minimally displaced set of an irreducible automorphism of a free group F_N . Our goal is to produce an essentially elementary proof that an irreducible automorphism acts co-compactly on its minimally displaced set (equivalently,

E-mail addresses: stefano.francaviglia@unibo.it, A.Martino@soton.ac.uk,
d.Syrigos@soton.ac.uk.

1991 *Mathematics Subject Classification.* 20E06, 20E36, 20E08.

the train track points) in Culler-Vogtmann space. We note that stronger results were produced by [15], who showed that the action on the *axis bundle* is co-compact, from which one easily deduces the result. However, their result only holds for non-geometric iwip automorphisms, and we believe our argument to be significantly simpler.

Irreducible automorphisms of F_N play a central role in the study of the outer automorphism group $Out(F_N)$, as they can be studied using the powerful train track machinery, which has been introduced by Bestvina and Handel in [3]. Also, they are generic elements of $Out(F_N)$ in the sense of random walks (see [20]).

Culler-Vogtmann Outer space CV_N is considered a classical method to understand automorphisms of free groups, by studying the action of $Out(F_N)$ on CV_N . More recently, there is an extensive study of the so-called Lipschitz metric on CV_N which seems that it has interesting applications (for example, see [1], [2], [11], [19]).

In particular, given an automorphism ϕ , we can define the displacement of ϕ in CV_N , as $\lambda_\phi = \inf\{\Lambda(X, (X)\phi) : X \in CV_N\}$ (where we denote by Λ the asymmetric Lipschitz metric). The set of minimally displaced points (or simply Min-Set) $Min(\phi)$ has interesting properties. For instance, the first two authors studied the Min-Set in [14] (in the context of general deformation spaces), where they proved that it is connected. As an application they gave solutions to some decision problems for irreducible automorphisms.

For an automorphism ϕ the centraliser, $C(\phi)$, preserves the Min-Set, $Min(\phi)$. Our main result is the following:

Theorem 3.1. *Let ϕ be an irreducible automorphism of F_N with $\lambda_\phi > 1$. The quotient space $Min(\phi)/C(\phi)$ is compact.*

As we have already mentioned, Handel and Mosher prove the previous theorem (under some extra hypotheses) in [15]. It is worth mentioning that some of the main ingredients of their proof is generalised by Bestvina, Guirardel and Horbez in the context of deformation spaces of free products (see [5]).

We then collect known results together, showing that the centraliser of an irreducible automorphism with $\lambda_\phi > 1$ is virtually cyclic, and deduce the following.

Theorem 3.3. *Let ϕ be an irreducible automorphism of F_N with $\lambda_\phi > 1$. Then $Min(\phi)/C(\phi)$ is compact.*

Remark. Note that centralisers for iwip automorphisms are well known to be virtually cyclic, but the more general statement is also true since irreducible automorphisms of infinite order which are not iwip are geometric, and their centralisers are also geometric, hence virtually cyclic. We collect these observations in more detail in the proof of Theorem 3.3.

In order to have a complete picture for irreducible automorphisms of a free group, we study irreducible automorphisms of growth rate one (all of these have finite order). In this case, we have the following:

Theorem 3.8. *Let ϕ be an irreducible automorphism of F_N with $\lambda_\phi = 1$. There is a single point $T \in CV_N$ so that $Min(\phi) = Fix(\phi) = \{T\}$.*

As an application, we get the following corollary:

Corollary 3.9. *Let ϕ be an irreducible automorphism of F_N with $\lambda_\phi = 1$. There is some $T \in CV_N$ so that $C(\phi)$ fixes T . In particular, $C(\phi)$ is finite.*

2. PRELIMINARIES

2.1. Culler -Vogtmann space. For the rest of the paper, we will denote by F_N the free group on N generators, for some $N \geq 2$. Firstly, we will describe the construction of the Culler-Vogtmann space which is denoted by CV_N and it is a space on which $Out(F_N)$ acts nicely.

Let's fix a free basis x_1, \dots, x_N of F_N . We denote by R_N the rose with n -petals where we identify each x_i with a single petal of R_N (the petals are still denoted by x_1, \dots, x_N).

Definition 2.1. A marked metric graph of rank n , is a triple (T, h, ℓ_T) so that :

- T is a graph (without valence one or two vertices) with fundamental group isomorphic to F_N .
- $h : R_N \rightarrow T$ is a homotopy equivalence, which is called the marking
- $\ell_T : E(T) \rightarrow (0, 1)$ is a metric on T , which assigns a positive number on each edge of T , with the property $\sum_e \ell_T(e) = 1$.

We are now in position to define Culler Vogtmann space:

Definition 2.2. CV_N is the space of equivalence classes, under \sim , of marked metric graphs of rank n . The equivalence relation \sim is given by $(T, h, \ell_T) \sim (S, h', \ell_S)$ if and only if there is an isometry $g : T \rightarrow S$ so that gh is homotopic to h' .

In order to simplify the notation, whenever ℓ_T, h are clear by the context, we can simply write T instead of the triple (T, h, ℓ_T) .

Remark. Equivalently, one may take universal covers to get a different formulation of Culler Vogtmann space; as the space of free, minimal, simplicial F_N trees of volume 1, up to equivariant isometry.

Action of $Out(F_N)$: Let $\phi \in Out(F_N)$ and (T, h, ℓ_T) be a marked metric graph. We define $((T, h, \ell_T))\phi = (T, h', \ell_T)$ where $h' = h\phi$ (here we still denote by ϕ , the natural representative of ϕ as a homotopy equivalence from R_N to R_N).

Simplicial Structure. If we fix a pair (T, h) of a topological marked graph (without metric) and we consider all the different possible metrics on $E(T)$ (i.e. assignments of a positive number on each edge so that the volume of T is one), we obtain an (open) simplex in CV_N with dimension $|E(T)| - 1$. However, if we allow some edges to have length 0, then it is not always true that the new graph will have rank n , so the resulting simplex is not necessarily in CV_N (note that some faces could be in CV_N). Therefore, CV_N is not a simplicial complex, but it can be described as a union of open simplices. Note that it follows immediately by the definitions, that $Out(F_N)$ acts simplicially on CV_N . In the following theorem, we list some more properties which are proved in [6].

Theorem (See [6]).

- i) CV_N is contractible.*
- ii) The maximum dimension of a simplex is $3N - 4$. In other words, the maximum number of edges in a marked metric graph with rank N , is bounded above by $3N - 3$.*
- iii) $Out(F_N)$ acts on CV_N by finite point stabilisers and the quotient space is consisted of finitely many open simplices*

Translation length. Given a marked metric graph T , we would like to define the translation length of the conjugacy class of a non-trivial group element a with respect to T . It is natural to define the length of a , as the sum of lengths of the edges that a crosses when it is realised as a reduced loop in T . We define the translation length of the

conjugacy class of $a \in F_N$, as $\ell_T([a]) = \inf\{\text{len}_T(a') : [a'] = [a]\}$. It is easy to see that any group element is freely homotopic to an embedded loop which realises the minimum.

Remark. This translation length is the same as the translation length - in the sense of the minimum distance moved by the group element - of the corresponding action of the group element on the universal cover, a tree.

Thick part. We can now define the thick part of CV_N , which will be essential in our arguments:

Definition 2.3. Let $\epsilon > 0$. We define the thick part of CV_N , and we denote it by $CV_N(\epsilon)$, the subspace of CV_N which is consisted of all the points $T \in CV_N$ with the property that every non-trivial conjugacy class α of F_N has translation length at least ϵ , i.e. $\ell_T(\alpha) \geq \epsilon$.

Centre of a simplex. We will need the notion of a special point of a simplex, the centre, so we give the following definition:

Definition 2.4. Let Δ be a simplex of CV_N . The point of Δ where all the edges have the same length is called the centre of Δ and we will denote it by X_Δ .

Remark 2.5. By the definition of the action of $Out(F_N)$ on CV_N , for any automorphism ψ , $X_{\psi(\Delta)} = \psi(X_\Delta)$, i.e. the centre of Δ is sent to the centre of $\psi(\Delta)$.

2.2. Stretching factor and Automorphisms. We define a natural notion of distance on CV_N , which has been studied in [10].

Definition 2.6. Let $T, S \in CV_N$. We define the (right) stretching factor as:

$$\Lambda(T, S) = \sup\left\{\frac{\ell_S([a])}{\ell_T([a])} : 1 \neq a \in F_N\right\}$$

In the following Proposition, we state some main properties of the stretching factor:

Proposition 2.7 (See [10]).

- i) For any two points $T, S \in CV_N$, $\Lambda(T, S) \geq 1$, with equality if and only if $T = S$.*
- ii) For any $T, S, Q \in CV_N$, the (non-symmetric) multiplicative triangle inequality for Λ holds. In other words, $\Lambda(T, S) \leq \Lambda(T, Q)\Lambda(Q, S)$.*
- iii) For any $\psi \in Out(F_N)$ and for any $T, S \in CV_N$, $\Lambda(T, S) = \Lambda((T)\psi, (S)\psi)$*

Note that Λ is not symmetric, i.e. there are points T, S of CV_N so that $\Lambda(T, S) \neq \Lambda(S, T)$. In fact, it is not even quasi-symmetric, in general. However, it is proved in [1] that for any $\epsilon > 0$, its restriction on the ϵ -thick part, $CV_N(\epsilon)$, induces a quasi-symmetric function. More specifically, we have the following:

Proposition 2.8 (Theorem 24, [1]). *For every $\epsilon > 0$ and for every $T, S \in CV_N(\epsilon)$, there is a uniform constant K (depending only on ϵ and N) so that $\Lambda(T, S) \leq \Lambda(S, T)^K$.*

One could consider the function $d_R(T, S) = \ln \Lambda(T, S)$, which behaves as an asymmetric distance, or even the symmetrised version $d(T, S) = d_R(T, S) + d_R(S, T)$, which is a metric on CV_N . We choose to work with Λ , as it is related more naturally to the displacement of an automorphism which is defined in the next subsection. However, as we will work only with some fixed ϵ -thick part of CV_N , it follows by the quasi-symmetry of Λ , that we could use exactly the same arguments using d , instead.

Balls in Outer Space. The asymmetry implies that there are three different types of balls that we can define, which are different, in general.

- (1) The symmetric closed ball with centre $T \in CV_N$ and radius $r > 0$:

$$B(T, r) = \{X \in CV_N : \Lambda(X, T)\Lambda(T, X) \leq r\}$$

(2) The in-going ball with centre $T \in CV_N$ and radius $r > 0$:

$$B_{in}(T, r) = \{X \in CV_N : \Lambda(X, T) \leq r\}$$

(3) The out-going ball with centre $T \in CV_N$ and radius $r > 0$:

$$B_{out}(T, r) = \{X \in CV_N : \Lambda(T, X) \leq r\}$$

Proposition 2.9. *Let $T \in CV_N$ and $r > 0$.*

- i) The symmetric ball $B(T, r)$ is compact.*
- ii) Then in-going ball $B_{in}(T, r)$ is compact.*

Proof.

- i) The statement is proved in [10], Theorem 4.12.
- ii) Let T, r as before. Firstly note that there is some C (the injective radius of T) for which $\ell_T([g]) \geq C > 0$ for every non-trivial $g \in F_N$. Therefore, for every non-trivial $g \in F_N$ and for every $X \in B_{in}(T, r)$, it holds that:

$$\frac{\ell_T([g])}{\ell_X([g])} \leq \Lambda(X, T) \leq r \Rightarrow \ell_X([g]) \geq \frac{\ell_T([g])}{r} \geq \frac{C}{r}$$

As a consequence, it follows that $B_{in}(T, r) \subset CV_N(C/r)$. By quasi-symmetry of Λ when it is restricted on the thick part (2.8), we have that there is some M so that

$$B_{in}(T, r) \subseteq B(T, r^{2M})$$

The result now follows by i), as $B_{in}(T, r)$ is a closed subset of the compact set $B(T, r^{2M})$ and therefore compact. □

Note that it's not difficult to find an out-going ball $B_{out}(T, r)$, for some $T \in CV_N$, $r > 0$, which is not compact. It is worth mentioning that B_{out} is weakly convex, while B_{in} is not, in general (as is proved in [19]).

In [10], it is proved that the supremum in the definition of stretching factor $\Lambda(T, S)$ is maximum, as there is a hyperbolic element that realises the supremum. Even more, such a hyperbolic element can be chosen by a finite list of candidates of T .

Definition 2.10. Let $T \in CV_N$. A hyperbolic element a of F_N is called a candidate with respect to T , if the realisation of a as a loop in T is either:

- an embedded circle
- a figure eight
- a barbell

The set of candidates with respect to T is called the set of candidates and it is denoted by $Cand(T)$.

Proposition 2.11 (Prop.3.15, [[10])). *For any $T, S \in CV_N$, there is a candidate $a \in Cand(T)$ so that:*

$$\Lambda(T, S) = \frac{\ell_S([a])}{\ell_T([a])}$$

Displacement. Now let's fix an automorphism $\phi \in Out(F_N)$. The displacement of ϕ is $\lambda_\phi = \inf\{\Lambda(X, \phi(X)) : X \in CV_N\}$.

Definition 2.12. Let ϕ be an outer automorphism of F_N . Then we define the Min-Set of ϕ , as $Min(\phi) = \{T \in CV_N : \Lambda(T, \phi(T)) = \lambda_\phi\}$.

Note that $Min(\phi)$ could be empty, in general. However, it is non-empty when ϕ is irreducible (see [12] for more details).

Definition 2.13. Let ϕ be an automorphism of F_N . Then ϕ is called reducible, if there is a free product decomposition of $F_N = A_1 * \dots * A_k * B$ (where B could be empty) so that every A_i , $i = 1, \dots, k$ is proper free factor and the conjugacy classes of A_i 's are permuted by ϕ . Otherwise, ϕ is called irreducible.

We note that by [Th.8.19, [12]], $Min(\phi)$ coincides with the set of train track points of ϕ , when ϕ is irreducible. That is, the points of CV_N that admit a (not necessarily simplicial) train track representative of ϕ .

2.3. Properties of the Thick Part. In this subsection, we list some properties of the thick part that we will need in the following section.

Proposition 2.14. *Let ϕ be an irreducible automorphism of F_N . Then there is a positive number ϵ_1 (depending only on N and on ϕ) so that $Min(\phi) \subset CV_N(\epsilon_1)$. One could take $\epsilon_1 = 1/((3N - 3)\mu^{3N-2})$ for any $\mu > \lambda_\phi$.*

Proof. For a proof see Proposition 10 of [2]. □

Lemma 2.15. *Let Δ be a simplex of CV_N and T, S be two points of Δ . If we further suppose that $T \in CV_N(\epsilon)$, then $\Lambda(T, S) \leq \frac{2}{\epsilon}$.*

Proof. The proof is an immediate corollary of the candidates theorem (2.11). Firstly, note that candidates depend only on the simplex and so for any $T, S \in \Delta$, $Cand(T) = Cand(S) = \mathcal{C}$. The result now follows by the remark that for every $g \in \mathcal{C}$ we have $\ell_S([g]) \leq 2$, while for every $1 \neq g$, $\ell_T([g]) \geq \epsilon$ as $T \in CV_N(\epsilon)$. □

Remark 2.16. Let Δ be any simplex of CV_N , then there is some uniform positive number ϵ_2 (depending only on N), so that the centre of Δ is ϵ_2 -thick. One could take, $\epsilon_2 = 1/(3N - 3)$. Therefore, it is easy to see $\epsilon_1 < \epsilon_2$ and so the centre of any simplex is ϵ_1 -thick.

Proof. The number of edges in any graph is bounded above by $3N - 3$. Therefore the length of each edge with respect to the centre of a simplex, is bounded below by $1/(3N - 3)$. The same number gives us an obvious lower bound for the translation length of any hyperbolic element. □

2.4. Connectivity of the Min-set. In this subsection, we will state some results from [13] and [14], that we will need in the following section.

Definition 2.17. Let $T, S \in CV_N$. A *simplicial path* between T, S is given by:

- (1) A finite sequence of points $T = X_0, X_1, \dots, X_k = S$, such that for every $i = 1, \dots, k$, there is a simplex Δ_i such that the simplices $\Delta_{X_{i-1}}$ and Δ_{X_i} are both faces of Δ_i .
- (2) Euclidean segments $\overline{X_{i-1}X_i}$.

The simplicial path is then the concatenation of these Euclidean segments.

The following results were proved in [13] and [14], in the more general context of deformation spaces of free products.

Theorem 2.18. *Let ϕ be an automorphism of F_N .*

- (1) *Let Δ be a simplex of CV_N . If $X, Y \in Min(\phi) \cap cl(\Delta)$, where $cl(\Delta)$ is the closure of Δ in CV_N . Then the Euclidean segment \overline{XY} is contained to $Min(\phi)$.*

- (2) If ϕ is irreducible automorphism of F_N , then $Min(\phi)$ is connected by simplicial paths in CV_N ; that is, for every $T, S \in Min(\phi)$ there is a simplicial path T and S , which is entirely contained in $Min(\phi)$.

Proof.

- (1) This is an immediate consequence of the quasi-convexity of the displacement function, see Lemma 6.2 in [13].
- (2) This follows by the Main Theorem of [14] (see Theorem 5.3.) which implies that the set of minimally displaced points for ϕ as a subset of the free splitting complex (i.e. the simplicial boardification of CV_N) of F_N is connected by simplicial paths, combined with the fact that the Min-Set of an irreducible automorphism does not enter some thin part of CV_N (see the previous subsection). It is easy to see now that, by the connectivity, it's not possible to have minimally displaced points in the boundary without entering any thin part. As a consequence, any minimally displaced point of the free splitting complex must be a point of CV_N and the result follows. □

3. RESULTS

3.1. Exponential growth. Firstly, we will concentrate on the case where ϕ is irreducible with exponential growth. In this case, we have the following theorem:

Theorem 3.1. *Let ϕ be an irreducible automorphism of F_N with $\lambda_\phi > 1$. The quotient space $Min(\phi)/C(\phi)$ is compact.*

Proof. We will prove that there is a compact set \mathcal{K} of CV_N so that $Min(\phi) \subseteq \mathcal{K}C(\phi)$ and the theorem follows, as $Min(\phi)$ is closed. Therefore as in-going balls are compact (by 2.9), the compact set \mathcal{K} can be chosen to be an in-going closed ball. Let's fix a point $T \in Min(\phi)$. It is then sufficient to prove that there is some positive radius L so that for any $X \in Min(\phi)$, there is some element α of $C(\phi)$ which satisfies $\Lambda(X, (T)\alpha) \leq L$.

We will argue by contradiction. Let's suppose that there is a sequence of points $X_m \in Min(\phi)$, $m = 1, 2, 3, \dots$, so that $\Lambda(X_m, (T)\alpha) \geq m$, $m = 1, 2, \dots$ and for every $\alpha \in C(\phi)$.

Note that there are finitely many $Out(F_N)$ - orbits of open simplices on CV_N . Therefore, up to taking a subsequence of X_m , we can suppose that there is an (open) simplex Δ and a sequence of (difference of markings) $\psi_m \in Out(F_N)$ so that $X_m \in (\Delta)\psi_m$.

Firstly, we apply Proposition 2.14 for ϕ and we get a constant ϵ (depending on N and on ϕ), so that $Min(\phi) \subset CV_N(\epsilon)$. Also, by the Remark 2.16, the centre of any simplex is ϵ -thick. In particular, the centre X_Δ of Δ belongs to $CV_N(\epsilon)$ and $(X_m)\psi_m \in \Delta$, so by 2.15 there is a constant $M = M(\epsilon)$ (which doesn't depend on Δ or m) which satisfies $\Lambda(X_\Delta, (X_m)\psi_m^{-1}) \leq M$. It follows that $\Lambda(X_m, (X_\Delta)\psi_m) \leq M$ for every m .

In addition, by assumption, $X_m \in Min(\phi)$ which is equivalent to $\Lambda(X_m, (X_m)\phi) = \lambda_\phi = \lambda$. Therefore, as an easy application of the multiplicative triangle inequality for Λ and the fact that $Out(F_N)$ acts on CV_N by isometries with respect to Λ , the previous relations imply that:

$$\begin{aligned} \Lambda((X_\Delta)\psi_m, (X_\Delta)\psi_m\phi) &\leq \Lambda((X_\Delta)\psi_m, X_m)\Lambda(X_m, (X_m)\phi)\Lambda((X_m)\phi, (X_\Delta)\psi_m\phi) \\ &= \Lambda(X_\Delta, (X_m)\psi_m^{-1})\Lambda(X_m, (X_m)\phi)\Lambda(X_m, (X_\Delta)\psi_m) \\ &\leq M^2\lambda. \end{aligned}$$

The previous inequality is equivalent to $\Lambda(X_\Delta, (X_\Delta)\psi_m\phi\psi_m^{-1}) \leq M^2\lambda$ for every m . Note that by the Remark 2.5, $(X_\Delta)\psi_m\phi\psi_m^{-1}$ are the centres of the corresponding simplices $(\Delta)\psi_m\phi\psi_m^{-1}$ for every m .

As CV_N is locally finite, there are finitely many simplices so that their centers have bounded distance from X_Δ . Therefore, it follows that infinitely many of the simplices $(\Delta)\psi_n\phi\psi_n^{-1}, n = 1, 2, \dots$ must be the same, which means that after possibly taking a subsequence of ψ_n , we have that $(X_\Delta)\psi_n\phi\psi_n^{-1} = (X_\Delta)\psi_m\phi\psi_m^{-1}$ for every n, m . As a consequence, the automorphisms $(\psi_n\phi\psi_n^{-1})(\psi_1\phi^{-1}\psi_1^{-1})$ fix X_Δ for every n .

On the other hand, the stabiliser of any point of CV_N is finite and so infinitely many of these automorphisms are forced to be the same or, in other words, after taking a subsequence, we can suppose that $\psi_m\phi\psi_m^{-1} = \psi_n\phi\psi_n^{-1}$ for every n, m . This is equivalent to $\psi_m^{-1}\psi_n \in C(\phi)$ for every n, m and in particular, by fixing one of the indices to be $m = 1$, we get that for every n there is some $\alpha_n \in C(\phi)$, so that $\psi_n = \psi_1\alpha_n$.

As a consequence, the Λ -distance from $(X_\Delta)\psi_n$ to $(T)\alpha_n$ does not depend on n , as

$$\Lambda((X_\Delta)\psi_n, (T)\alpha_n) = \Lambda((X_\Delta)\psi_1\alpha_n, (T)\alpha_n) = \Lambda((X_\Delta)\psi_1, T) = C.$$

Note that as we have already seen $\Lambda(X_n, (X_\Delta)\psi_n) \leq M$ for every n . Therefore, by applying again the triangle inequality, it follows that $\Lambda(X_n, (T)\alpha_n) \leq MC$ for every n . This contradicts our assumption that $\Lambda(X_n, (T)\alpha_n) \geq n$, as neither M nor C depend on n (note that here by an abuse of the notation, where we still denote by X_n a subsequence X_{k_n} of the original X_n , but the inequality still holds because $k_n \geq n$ for every n). \square

Remark 3.2. In the previous proof we used irreducibility only in order to ensure the condition that there is some uniform ϵ so that $Min(\phi) \subset CV_N(\epsilon)$ which follows by the Proposition 2.14. Therefore, we could replace the assumption of irreducibility with this weaker condition.

For the centralisers of irreducible automorphisms with exponential growth, the following theorem holds:

Theorem 3.3. *Let ϕ be an irreducible automorphism of F_N with $\lambda_\phi > 1$. Then $Min(\phi) / \langle \phi \rangle$ is compact.*

Proof. In light of the previous theorem, it is sufficient to prove that the centraliser of an irreducible automorphism of exponential growth rate is virtually cyclic.

The case when ϕ is irreducible with irreducible powers (iwip) this is well known, by the main result of [4]. Note that if ϕ is atoroidal and irreducible with $\lambda_\phi > 1$, it follows by [16] that ϕ is iwip.

As a consequence, we suppose for the rest of the proof that ϕ is toroidal and irreducible. In this case, ϕ is a geometric automorphism, i.e. it is induced by a pseudo-Anosov automorphism f of a surface Σ with $p \geq 1$ punctures, which acts transitively on the boundary components (this was a folk theorem, until recently where the details appeared in an appendix of [18]). Note that ϕ is iwip exactly when $p = 1$.

It is well known (see [17]) that the centraliser $C_{MCG(\Sigma)}(f)$ of the pseudo-Anosov f in $MCG(\Sigma)$ is virtually cyclic. Therefore, it is enough to prove that the centraliser $C(\phi) = C_{Out(F_n)}(\phi)$ of ϕ in $Out(F_n)$ is isomorphic to $C_{MCG(\Sigma)}(f)$.

Let's denote by c_1, \dots, c_p the elements corresponding to the peripheral curves (a simple curve around each puncture). We also denote by $Out^*(F_n)$, the subgroup of the automorphisms that preserve the set of conjugacy classes of simple peripheral curves (which are c_i 's and their inverses). By the Dehn-Nielsen-Baer Theorem for surfaces with punctures (see Theorem 8.8. of [9]), we have that the natural map from $MCG(\Sigma)$ to $Out^*(F_n)$ is an isomorphism. In other words, an automorphism ψ of $Out(F_n)$ is induced by an element of

$MCG(\Sigma)$, exactly when it preserves the set of conjugacy classes of the peripheral curves, or equivalently $\psi \in Out^*(F_n)$.

We will now show that any element of $C(\phi)$ is induced by an element of $C_{MCG(\Sigma)}(f)$ and the proof follows, as any element of $C_{MCG(\Sigma)}(f)$ induces an element of $C(\phi)$. We can assume that without loss, after re-numbering if needed, that $[\phi(c_i)] = [c_{i+1}] \pmod p$. If $\psi \in C(\phi)$, then for every i , $[\phi\psi(c_i)] = [\psi\phi(c_i)] = [\psi(c_{i+1})] \pmod p$. It follows that f preserves the closed curves $[\psi(c_i)]$ and since f is pseudo-Anosov, we get that any such curve must be a peripheral curve, up to orientation (proper powers can be discounted as ψ is an automorphism). Therefore, ψ must preserve the set of (conjugacy classes) of the peripheral curves and so by the previous criterion, it is induced by an element of $MCG(\Sigma)$. □

As an immediate application of the previous two theorems, we get the following statement which seems to be known to the experts, but it doesn't appear explicitly in the literature:

Theorem 3.4. *Let ϕ be an irreducible automorphism of F_N with $\lambda_\phi > 1$. It holds that $Min(\phi)/\langle \phi \rangle$ is compact.*

3.2. Growth rate one. In this subsection, we will cover the case of irreducible automorphisms of growth rate one (it is known that they have finite order). This class has been studied by Dicks and Ventura in [8], where they give an explicit description of any such automorphism.

Notation 3.5. We describe below two types of topological graphs and a graph map for each case:

- For the first type, let p be an odd prime. In this case, we define a graph X_p which has rank $p - 1$. This graph is consisted of two vertices and p edges which connect them. In order to describe the graph map is more convenient to identify the set of vertices $\{v_0, v_1\}$ with the set $\{0, 1\}$ and the edge set $\{e_1, \dots, e_p\}$ with \mathbb{Z}_p , while all edges are oriented so that their initial vertex is v_0 and their terminal vertex v_1 . We denote by α_p the graph map, which fixes the vertices and sends e_i to $e_{i+1} \pmod p$.
- For the second type, let p, q two primes where $p < q$ (here we don't assume that p is odd). We define a graph X_{pq} of rank $pq - p - q + 1$ which is consisted of $p+q$ vertices and pq edges. More specifically, the set of vertices is consisted of two distinct subsets $\{v_1, \dots, v_p\}, \{w_1, \dots, w_q\}$ which can be naturally identified with $\mathbb{Z}_p \sqcup \mathbb{Z}_q$. Also, for every $i \in \mathbb{Z}_p$ and for every $j \in \mathbb{Z}_q$, there is a unique edge $e_{i,j}$ with initial vertex v_i and terminal vertex w_j and so we can naturally identify the set of edges with $\mathbb{Z}_p \times \mathbb{Z}_q$. We denote by α_{pq} the map which sends $e_{i,j}$ to $e_{i+1, j+1} \pmod p, \pmod q$, respectively).

Note that X_p , for an odd prime p , can be seen as an element of CV_{p-1} by assigning length $1/(p-1)$ on each edge and by considering a marking $R_{p-1} \rightarrow X_p$. Similarly, X_{pq} can be seen as an element of $CV_{pq-p-q+1}$. This is true even when $p = 2$, as in this case $X_{2q} = X_q$ (as elements of CV_{q-1}). We consider this case separately, as the maps α_q and α_{2q} are different (α_{2q} inverts the orientation of each edge, while α_q preserves it).

We can now state the main result of [8] (it is proved in Proposition 3.6, even if it is stated in a different form, the following formulation is evident by the proof):

Proposition 3.6. *If ϕ is an irreducible automorphism of F_N with $\lambda_\phi = 1$, then it can be represented by either α_p for some odd prime p or α_{pq} for some primes $p < q$, as in Notation 3.5.*

We need the following lemma:

Lemma 3.7. *Let ϕ be an irreducible automorphism of CV_N with $\lambda_\phi = 1$. Let S be a point of $Min(\phi)$ and $f : S \rightarrow S$ be an optimal representative of ϕ . Then f is an isometry on S , and hence a graph automorphism. Moreover, f is the unique optimal map representing ϕ on S .*

Proof. Note that as $\lambda_\phi = 1$, $Min(\phi)$ is simply the set of fixed points of ϕ in CV_N . Hence for any loop, γ , the lengths of γ and $f(\gamma)$ are the same (as loops) in S , and from there it follows easily that f is an isometry.

We can lift this isometry to the universal cover and invoke [7], to conclude that this isometry is unique (up to a covering translation), and hence all optimal maps are the same on $Min(\phi)$.

Alternatively, notice that two graph automorphisms give rise to the same action on the associated simplex if and only if the graph maps are the same. \square

Remark. Note that this uniqueness statement is definitely false when $\lambda_\phi > 1$; see [13], Example 3.14.

We can now prove our result for the finite order irreducible automorphisms of F_N .

Theorem 3.8. *Let ϕ be an irreducible automorphism of F_N with $\lambda_\phi = 1$. There is a single point $T \in CV_N$ so that $Min(\phi) = Fix(\phi) = \{T\}$.*

Note that the existence of such a point T is proved in [8]; the content of this Theorem is the uniqueness of such a point.

Proof. As note above, the fact that $\lambda_\phi = 1$ implies that $Min(\phi) = Fix(\phi)$.

By applying Proposition 3.6, we get that ϕ can be represented by either α_p or α_{pq} (for some primes p, q) as an isometry of X_p or X_{pq} (as in notation 3.5), respectively. For the rest of the proof we will denote the graph map by α and the graph by T . In particular, $\Delta_T = \Delta$, then ϕ fixes the centre (i.e. if we assign the same length to every edge of T) of Δ , as an element of CV_N . We will prove that T (with the metric given as above) is the unique fixed point of CV_N for ϕ .

By the second assertion of Theorem 2.18, $Fix(\phi) = Min(\phi)$ is connected by simplicial paths in CV_N . Now consider some other point, $S \in CV_N$ fixed by ϕ . If we connect S and T by a (simplicial) path in $Fix(\phi)$, we will be able to produce a point $T' \in Fix(\phi)$ such that either Δ' is a face of Δ or vice versa (where Δ' is the simplex defined by T').

However, it is clear that ϕ cannot fix any other point of Δ ; this is because α acts as a cyclic permutation of the edges of T , and hence the only metric structure that can be preserved assigns the same length to every edge. Therefore, Δ must be a face of Δ' .

We will aim to show that Δ must be equal to Δ' , which will prove our result. (Since then the only possibility left is that $S = T$.)

Since Δ is a face of Δ' , there exists a forest, F , whose components are collapsed to produce T from T' (as graphs, absent the metric structure). However, the connectivity of $Fix(\phi)$ allows us to connect T to T' - as metric graphs - via a Euclidean segment in the closure of Δ' (see the first assertion of 2.18). Thus, without loss of generality, we may assume that the optimal map representing ϕ on T' - call this α' - leaves F invariant, since α' is an isometry (by Lemma 3.7), and so if the volume of F is sufficiently small, it must be sent to itself.

Our goal is to show that, under the assumption that T' has no valence one or two vertices, each component of F is a vertex. Hence $T = T'$.

Now if we ignore the metric structures we get that on collapsing F , α' induces a graph map on the quotient, which is T (as a graph). Since this must also represent ϕ , we deduce that this induced map is equal to α . (Alternatively, collapse F and assign the same length to the surviving edges. The map α' induces an isometry of this graph which is equal to α , by 3.7.)

Let G denote the cyclic group generated by α , acting on T . By the comments above, G has an action on T' so that collapsing components of F gives rise to the original action on T .

For the remainder of the proof, the edges of F will be called *black*, the edges of the complement of F will be called *white* and vertices that are incident to both and white edges will be called *mixed*. Accordingly, vertices have *black* valence and *white* valence respectively.

Now consider a component C of F . Note that we can think of C as a *vertex* of T . Let v_1, \dots, v_k be the leaves of C ; that is, those vertices of C with black valence equal to 1. Since T' can admit no valence one vertices (if we count both black and white edges), each v_i is mixed, incident to both black and white edges. Let ∂C denote the boundary of C in T' ; that is, the edges (necessarily white) of T' connecting some v_i to a vertex not in C .

Since a vertex stabiliser of T in G acts freely and transitively on the edges incident to it, we deduce that $Stab(C)$ (the set-wise stabiliser) acts freely and transitively on ∂C . Note that the transitive action on ∂C implies that $Stab(C)$ must also act transitively on the leaves, v_1, \dots, v_k .

Now $Stab(C)$ has prime order (since vertex stabilisers in T have prime order), so using the orbit-stabiliser theorem, we deduce that either $k = 1$ or $Stab(C)$ acts freely and transitively on v_1, \dots, v_k . In the former case, C consists of a single vertex. In the latter case we get that $k = |\partial C| = |Stab(C)|$. This implies that the white valence of each v_i is equal to 1. But now the valence of each v_i is exactly 2, leading to our desired contradiction.

Note that this argument doesn't quite work in the cases where T has a vertex of valence 2 - in the second case of Notation 3.5 when $p = 2$. Here we get vertices of valence 2 since the graph map α acts as a cyclic permutation on the edges along with an inversion, and we subdivide at the midpoints of edges which are fixed.

But in this case, the valence 2 vertices are a notational convenience, and we can omit them from T , and reach the same conclusion with the same argument. □

The following corollary is now immediate:

Corollary 3.9. *Let ϕ be an irreducible automorphism of F_N with $\lambda_\phi = 1$. There is some $T \in CV_N$ so that $C(\phi)$ fixes T . In particular, $C(\phi)$ is finite.*

Proof. It follows immediately by the previous Theorem, by noting that $Min(\phi)$ is $C(\phi)$ -invariant. □

In fact, the graph T in the previous corollary, is some graph as in Notation 3.5, we can get a much more precise description of the centraliser in each case.

Corollary 3.10. *If ϕ is irreducible automorphism of F_N of growth rate one, then $C(\phi)$ fixes a point X , where X is as in Notation 3.5. As a consequence, $C(\phi) = \langle \phi \rangle \times \langle \sigma \rangle$, where σ is the order two automorphism of F_N that is induced by the graph map of X sending every edge to its inverse.*

REFERENCES

- [1] Y. Algom-Kfir and M. Bestvina, *Geom Dedicata* (2012) 156: 81. <https://doi.org/10.1007/s10711-011-9591-2>
- [2] M. Bestvina, *A Bers-like proof of the existence of train tracks for free group automorphisms*, *Fund. Math.* **214** (2011), no. 1, 1–12.
- [3] M. Bestvina and M. Handel. *Train Tracks and Automorphisms of Free Groups.*, *Annals of Mathematics*, vol. 135, no. 1, 1992, pp. 151. JSTOR, www.jstor.org/stable/2946562.
- [4] M. Bestvina, M. Feighn, M. and Handel (1997). *Laminations, trees, and irreducible automorphisms of free groups*. *Geometric and Functional Analysis GAFA*, 7(2), 215–244.
- [5] M. Bestvina, V. Guirardel and C. Horbez (2017) *Boundary amenability of $Out(F_N)$* , arXiv:1705.07017
- [6] M. Culler and K. Vogtmann, *Moduli of graphs and automorphisms of free groups*, *Invent. Math.* **84** (1986), no. 1, 91–119. MR 830040
- [7] Marc Culler and John W. Morgan. Group actions on \mathbf{R} -trees. *Proc. London Math. Soc.* (3), 55(3):571–604, 1987.
- [8] W. Dicks and E. Ventura, *Irreducible automorphisms of growth rate one*, *Journal of Pure and Applied Algebra*, Volume 88, Issues 13,1993, Pages 51-62,ISSN 0022-4049, [https://doi.org/10.1016/0022-4049\(93\)90012-I](https://doi.org/10.1016/0022-4049(93)90012-I).
- [9] B. Farb and D. Margalit, *A primer on mapping class groups*, Princeton University press, 2012
- [10] S. Francaviglia and A. Martino, *Metric properties of outer space*, *Publ. Mat.* **55** (2011), no. 2, 433–473. MR 2839451
- [11] ———, *The isometry group of outer space*, *Adv. Math.* **231** (2012), no. 3-4, 1940–1973. MR 2964629
- [12] S. Francaviglia and A. Martino, *Stretching factors, metrics and train tracks for free products*. *Illinois J. Math.* 59 (2015), no. 4, 859–899. doi:10.1215/ijm/1488186013. <https://projecteuclid.org/euclid.ijm/1488186013>
- [13] ———, *Displacements of automorphisms of free groups I: Displacement functions, minpoints and train tracks*, preprint, arXiv1807.02781.
- [14] ———, *Displacements of automorphisms of free groups II: Connectedness of level sets*, preprint, arXiv:1807.02782.
- [15] M. Handel and L. Mosher, *Axes in outer space*”, American Mathematical Society Providence, RI, *Memoirs of the American Mathematical Society*, 2012, <http://cds.cern.ch/record/2122547>,
- [16] I. Kapovich, *Algorithmic detectability of iwip automorphisms*. *Bulletin of the London Mathematical Society* 46 (2014), no. 2, 279290.
- [17] J. McCarthy, *Normalizers and centralizers of pseudo-Anosov mapping classes*, preprint, June 8 (1994).
- [18] J.P. Mutanguha, *Irreducibility of a Free Group Endomorphism is a Mapping Torus Invariant*, preprint, arXiv: arXiv:1910.04285 (2019)
- [19] Y. Qing and K. Rafi. *Convexity of balls in the outer space.*, preprint, arXiv:1708.04921 (2017).
- [20] I. Rivin, *Walks on groups, counting reducible matrices, polynomials, and surface and free group automorphisms*. *Duke Math. J.* 142 (2008), no. 2, 353–379. doi:10.1215/00127094-2008-009. <https://projecteuclid.org/euclid.dmj/1206642158>